
Interner Bericht

Variance reduction by means of deterministic
computation: Collision estimate

Stefan Heinrich

285/96

Fachbereich Informatik

Universität Kaiserslautern · Postfach 3049 · D-67653 Kaiserslautern

Variance reduction by means of deterministic
computation: Collision estimate

Stefan Heinrich

285/96

Universität Kaiserslautern
AG Numerische Algorithmen
Postfach 30 49
67653 Kaiserslautern
Germany

Okt. 1996

Herausgeber: AG Numerische Algorithmen
Leiter: Professor Dr. S. Heinrich

Variance reduction by means of deterministic computation: Collision estimate

Stefan Heinrich

Universität Kaiserslautern
Fachbereich Informatik
Postfach 30 49
67653 Kaiserslautern
email: heinrich@informatik.uni-kl.de

Abstract

We study the collision estimate of Monte Carlo methods for the solution of integral equations. A new variance technique is proposed and analyzed. It consists in the separation of the main part by constructing a neighboring equation based on deterministic numerical methods.

Keywords: Monte Carlo method, variance reduction, collision estimate.

1991 *Mathematics Subject Classification*. Primary 65C05

1 Introduction

In a recent paper, [Hei95], a new variance reduction technique was introduced for the Monte Carlo solution of Fredholm integral equations. The idea, based on work in complexity theory, [HM93], [Hei94], consists in constructing a new equation sufficiently close to the original one and then applying standard schemes to both equations simultaneously. So the approach is a special case of the separation of main part (also called control variate) technique. As shown in [Hei95], neighboring equations can be constructed by exploiting the system and the approximate solution of deterministic schemes of solving the equation. The gain in variance reduction can be controlled by the discretization error. Hence, by applying the Monte Carlo method with n samples, the overall error is essentially the deterministic discretization error multiplied by the classical Monte Carlo rate $n^{-1/2}$ (see [HM93], [Hei94] for a theoretical foundation of this statement). Considerable improvements are possible this way as experiments in [Hei95] showed.

In [Hei95], exact expressions for the variance of the method as well as estimates in terms of proximity of the two equations were obtained for one of the classical Monte

Carlo algorithms - the absorption scheme. As a rule (compare [Erm71], [Mik91a]), this is the technically simpler case. The question arises what happens for the other classical method - the collision scheme. This is the theme of the present paper. We present and analyze the new technique for the collision scheme and prove that the variance is dominated by the square of the proximity of the respective kernels and right hand sides in some function space norms. This means the results of [Hei95] carry over to the collision estimate. The analysis is different and more complicated, but will be based on the results of [Hei95], and we shall work in a similar framework. Nevertheless, we try to keep the present paper selfcontained by recalling all needed notions and results.

Once the variance analysis of [Hei95] is extended to the collision estimate, we can apply the other results of that paper: The Galerkin method can be used to construct a neighboring equation and the proximity of kernels and right-hand sides can be estimated from certain parameters of that method. We do not repeat this here, but refer to [Hei95] and [Hei96] instead.

General references for Monte Carlo methods are [SG69], [Erm71], [Sob73], [EM82], [KW86], [ENS89], [Mik91a], [Mik91b], [Sab91]. Predecessors of our approach can be found in [Erm71], ch. 6.2.5, [Spa79], [ES85], [Mik91b], §5.10, [Sab91], ch. 2.2.3, [LW94].

2 The Algorithm

We consider the Fredholm integral equation of the second kind

$$u(x) = \int_X K(x, y)u(y) d\mu(y) + f(x). \quad (1)$$

Here X is a non-empty set, endowed with a σ -algebra Σ of subsets and a finite positive, σ -additive measure μ on (X, Σ) . If $1 \leq s < \infty$, $L_s(X) = L_s(X, \Sigma, \mu)$ stands for the space of s -integrable functions and if $s = \infty$ for the space of essentially bounded functions. We assume that $f \in L_\infty(X)$ and that K is $\Sigma \times \Sigma$ measurable and satisfies

$$\|K\|_{L_\infty(L_1)} := \operatorname{ess\,sup}_{x \in X} \int_X |K(x, y)| d\mu(y) < \infty.$$

This is equivalent to saying that the integral operator T_k defined for $g \in L_\infty(X)$ by

$$(T_K g)(x) = \int_X K(x, y)g(y) d\mu(y)$$

acts continuously in $L_\infty(X)$. Moreover,

$$\|T_K : L_\infty(X) \rightarrow L_\infty(X)\| = \|K\|_{L_\infty(L_1)}.$$

Our aim is to compute the value (u, Φ) of a functional $\Phi \in L_1(X)$ at the solution u of (1). Let $p_0(x)$ and $p(x, y)$ be non-negative measurable functions on (X, Σ) and $(X \times X, \Sigma \times \Sigma)$, respectively, satisfying

$$\int_X p_0(x) d\mu(x) = 1$$

and

$$\int_X p(x, y) d\mu(y) \leq 1 \quad (x \in X).$$

We shall assume that

$$\begin{aligned} \mu\{\Phi(x) \neq 0 \text{ and } p_0(x) = 0\} &= 0 \\ \mu \times \mu\{K(x, y) \neq 0 \text{ and } p(x, y) = 0\} &= 0. \end{aligned}$$

Define φ and k by

$$\begin{aligned} \Phi(x) &= \varphi(x)p_0(x) \\ K(x, y) &= k(x, y)p(x, y). \end{aligned}$$

We consider an absorbing Markov chain on X with density of initial distribution $p_0(x)$ and density of probability $p(x, y)$ of transition from x to y . We assume that the spectral radius of T_p in $L_\infty(X)$ is less than 1, hence almost all trajectories of the Markov chain are of finite length. Let

$$\xi = (x_0, x_1, \dots, x_m)$$

be such a trajectory. The classical collision estimate of the von Neumann Ulam scheme is defined as

$$\eta(k, f, \xi) = \sum_{\ell=0}^m \varphi(x_0)k(x_0, x_1) \dots k(x_{\ell-1}, x_\ell)f(x_\ell), \quad (2)$$

and it is well-known that

$$\mathbb{E}\eta(k, f, \xi) = \left((I - T_{kp})^{-1} f, \Phi \right) = (u, \Phi). \quad (3)$$

The approximation to (u, Φ) is then obtained by averaging N independent realizations of η . Given another pair of functions (h, g) (assumed to be close to (k, f) , see below), such that the exact solution v of

$$v = T_{hp}v + g$$

is known, we define a new random variable

$$\zeta(k, f, h, g, \xi) = (v, \Phi) + \eta(k, f, \xi) - \eta(h, g, \xi). \quad (4)$$

Observe that

$$\mathbb{E}\zeta(k, f, h, g, \xi) = (v, \Phi) + \left((I - T_{kp})^{-1} f, \Phi \right) - \left((I - T_{hp})^{-1} g, \Phi \right) = (u, \Phi).$$

Averaging over N independent trajectories, we approximate (u, Φ) by

$$\frac{1}{N} \sum_{i=1}^N \zeta(k, f, h, g, \xi_i) = (v, \Phi) + \frac{1}{N} \sum_{i=1}^N (\eta(k, f, \xi_i) - \eta(h, g, \xi_i)). \quad (5)$$

Hence $\eta(h, g, \xi)$ serves as the main part (control variate). The quality of the new scheme is determined by the variance of ζ :

$$\mathbb{E} \left((u, \Phi) - \frac{1}{N} \sum_{i=1}^N \zeta(k, f, h, g, \xi_i) \right)^2 = \frac{\text{Var}(\zeta)}{N}.$$

In the sequel we shall study this variance. As in [Hei95], we shall assume that

$$\varphi^2 p_0 \in L_1(X)$$

and that the functions k, h, f, g belong to the following classes: Fix $\alpha > 0, 0 < \gamma < 1, n_0 \in \mathbf{N}$ and define $\mathcal{K}(\alpha, \gamma, n_0)$ to be the set of all $k \in L_\infty(X^2)$ with

$$\|k\|_{L_\infty(X^2)} \leq \alpha$$

and

$$\|T_{k^2p}^{n_0} : L_\infty(X) \rightarrow L_\infty(X)\| \leq \gamma.$$

For $\theta > 0$ we let

$$\mathcal{F}(\theta) = \{f \in L_\infty(X); \|f\|_{L_\infty(X)} \leq \theta\}.$$

For $k, h \in \mathcal{K}(\alpha, \gamma, n_0)$ and $f, g \in \mathcal{F}(\theta)$ the collision estimate η and the new scheme ζ are well-defined random variables with finite second moment. (This is well-known, compare also Lemma 1 below, which we recall from [Hei95] for the sake of completeness.

Lemma 1 *Let $\alpha > 0$, $0 < \gamma < 1$, $n_0 \in \mathbf{N}$. If $k \in \mathcal{K}(\alpha, \gamma, n_0)$, then*

(i) $I - T_{k^2p}$ is invertible in $L_\infty(X)$ and $\|(I - T_{k^2p})^{-1} : L_\infty(X) \rightarrow L_\infty(X)\| \leq \beta_0$,
where $\beta_0 = (1 - \gamma)^{-1} \sum_{j=0}^{n_0-1} \alpha^{2j}$,

(ii) $\|kp\|_{L_\infty(L_1)} \leq \|k^2p\|_{L_\infty(L_1)}^{1/2} \leq \alpha$,

(iii) $\|T_{kp}^{n_0} : L_\infty(X) \rightarrow L_\infty(X)\| \leq \gamma$,

(iv) $I - T_{kp}$ is invertible in $L_\infty(X)$ and $\|(I - T_{kp})^{-1} : L_\infty(X) \rightarrow L_\infty(X)\| \leq \beta_1$,
where $\beta_1 = (1 - \gamma)^{-1} \sum_{j=0}^{n_0-1} \alpha^j$.

If $k, h \in \mathcal{K}(\alpha, \gamma, n_0)$, then

(v) $\|T_{khp}^{n_0} : L_\infty(X) \rightarrow L_\infty(X)\| \leq \gamma$, and $\|(I - T_{khp})^{-1} : L_\infty(X) \rightarrow L_\infty(X)\| \leq \beta_0$.

3 Variance of the new scheme

The following proposition provides a complicated, but exact expression for the variance of the new estimators ζ . Later on, we shall derive simpler forms yielding upper bounds for the variance.

Proposition 2 *Let $\alpha, \theta > 0$, $0 < \gamma < 1$, $n_0 \in \mathbf{N}$. Suppose that $k, h \in \mathcal{K}(\alpha, \gamma, n_0)$ and $f, g \in \mathcal{F}(\theta)$. Then the variance of $\zeta(k, f, h, g, \xi)$ can be expressed by the following formula:*

$$\begin{aligned}
\text{Var}(\zeta) = & \left((I - T_{k^2p})^{-1}(2f(I - T_{kp})^{-1}f - f^2) \right. \\
& - (I - T_{khp})^{-1}(2f(I - T_{hp})^{-1}g - fg) \\
& - (I - T_{khp})^{-1}(2g(I - T_{kp})^{-1}f - fg) \\
& + (I - T_{h^2p})^{-1}(2g(I - T_{hp})^{-1}g - g^2), \varphi^2 p_0 \Big) \\
& - \left((I - T_{kp})^{-1}f - (I - T_{hp})^{-1}g, \varphi p_0 \right)^2.
\end{aligned}$$

Proof: We represent the random variable $\zeta - (v, \Phi)$ by an infinite sum of random variables

$$\zeta - (v, \Phi) = \eta(k, f, \xi) - \eta(h, g, \xi) = \sum_{n=0}^{\infty} \zeta_n \quad (6)$$

where

$$\zeta_n = \varphi(x_0)(k(x_0, x_1) \cdots k(x_{n-1}, x_n)f(x_n) - h(x_0, x_1) \cdots h(x_{n-1}, x_n)g(x_n))$$

if $n \leq m(\xi)$ (the length of ξ) and $\zeta_n \cong 0$ if $n > m(\xi)$. Hence the sum (6) is in fact almost surely finite. For arbitrary $m, n \in \mathbb{N}$ with $n \geq m$ we have

$$\begin{aligned}
& \mathbb{E}\zeta_m\zeta_n \\
&= \int_{X^{n+1}} \varphi(x_0) (k(x_0, x_1) \cdots k(x_{m-1}, x_m)f(x_m) - h(x_0, x_1) \cdots h(x_{m-1}, x_m)g(x_m)) \\
&\quad \times \varphi(x_0) (k(x_0, x_1) \cdots k(x_{n-1}, x_n)f(x_n) - h(x_0, x_1) \cdots h(x_{n-1}, x_n)g(x_n)) \\
&\quad \times p(x_0)p(x_0, x_1) \cdots p(x_{n-1}, x_n) d\mu(x_0) \cdots d\mu(x_n) \\
&= (T_{k^2p}^m(fT_{kp}^{n-m}f) - T_{khp}^m(fT_{hp}^{n-m}g) - T_{hkp}^m(gT_{kp}^{n-m}f) + T_{h^2p}^m(gT_{hp}^{n-m}g), \varphi^2 p_0). \quad (7)
\end{aligned}$$

From (6) we get

$$\mathbb{E} \left(\sum_{n=0}^{\infty} \zeta_n \right)^2 = \mathbb{E} \sum_{m,n=0}^{\infty} \zeta_m\zeta_n = 2 \sum_{m=0}^{\infty} \sum_{n=m}^{\infty} \mathbb{E}\zeta_m\zeta_n - \sum_{m=0}^{\infty} \mathbb{E}\zeta_m^2, \quad (8)$$

where the operations on infinite series are justified, since the series (6) is absolutely convergent in the square mean (i.e. in the norm of L_2 over the probability space of the Markov chain). This follows from (7), the assumptions of the theorem and Lemma 1.

Now we combine (7) and (8). We apply the summation of (8) to each summand of (7). For the first one, we obtain

$$\sum_{m=0}^{\infty} \sum_{n=m}^{\infty} T_{k^2p}^m (f T_{kp}^{n-m} f) = \sum_{m=0}^{\infty} T_{k^2p}^m \left(f \sum_{\ell=0}^{\infty} T_{kp}^{\ell} f \right) = (I - T_{k^2p})^{-1} (f (I - T_{kp})^{-1} f)$$

and

$$\sum_{m=0}^{\infty} T_{k^2p}^m (f^2) = (I - T_{k^2p})^{-1} (f^2).$$

The remaining terms of (7) can be handled analogously. Hence we get

$$\begin{aligned} \mathbf{E}(\eta(k, f, \xi) - \eta(h, g, \xi))^2 &= ((I - T_{k^2p})^{-1} (2f(I - T_{kp})^{-1} f - f^2) \\ &\quad - (I - T_{khp})^{-1} (2f(I - T_{hp})^{-1} g - fg) \\ &\quad - (I - T_{khp})^{-1} (2g(I - T_{kp})^{-1} f - fg) \\ &\quad + (I - T_{h^2p})^{-1} (2g(I - T_{hp})^{-1} g - g^2), \varphi^2 p_0) \end{aligned} \quad (9)$$

which together with (6) and (3) proves the proposition.

Next we recall some notation from [Hei95], which we need in the sequel:

$$\begin{aligned} \delta(k, h) &= \|T_{kp} - T_{hp} : L_{\infty}(X) \rightarrow L_{\infty}(X)\| \\ &= \operatorname{ess\,sup}_{x \in X} \int_X |k(x, y) - h(x, y)| p(x, y) \, d\mu(y) \\ \varepsilon(k, h)^2 &= \|T_{(k-h)^2p} : L_{\infty}(X) \rightarrow L_{\infty}(X)\| \\ &= \operatorname{ess\,sup}_{x \in X} \int_X (k(x, y) - h(x, y))^2 p(x, y) \, d\mu(y). \end{aligned}$$

Hölder's inequality implies

$$\delta(k, h) \leq \varepsilon(k, h).$$

Theorem 3 *Let $\alpha, \theta > 0$, $0 < \gamma < 1$, and $n \in \mathbf{N}$. Then there exists a constant $c > 0$ such that for all $k, h \in \mathcal{K}(\alpha, \gamma, n_0)$ and $f, g \in \mathcal{F}(\theta)$*

$$\operatorname{Var}(\zeta(k, f, h, g, \xi)) \leq c(\varepsilon(k, h)^2 + \|f - g\|_{L_{\infty}(X)}^2).$$

Proof: We have according to (4) and (9)

$$\begin{aligned}
\text{Var}(\zeta) &\leq \mathbf{E}(\eta(k, f, \xi) - \eta(h, g, \xi))^2 \\
&\leq 2 \mathbf{E}(\eta(k, f, \xi) - \eta(h, f, \xi))^2 + 2 \mathbf{E}(\eta(h, f, \xi) - \eta(h, g, \xi))^2 \\
&= 2(w_1 + w_2, \varphi^2 p_0)
\end{aligned} \tag{10}$$

with

$$\begin{aligned}
w_1 &= (I - T_{k^2p})^{-1}(2f(I - T_{kp})^{-1}f - f^2) \\
&\quad - (I - T_{khp})^{-1}(2f(I - T_{hp})^{-1}f - f^2) \\
&\quad - (I - T_{khp})^{-1}(2f(I - T_{kp})^{-1}f - f^2) \\
&\quad + (I - T_{h^2p})^{-1}(2f(I - T_{hp})^{-1}f - f^2)
\end{aligned} \tag{11}$$

and

$$\begin{aligned}
w_2 &= (I - T_{h^2p})^{-1}(2f(I - T_{hp})^{-1}f - f^2) \\
&\quad - (2f(I - T_{hp})^{-1}g - fg) \\
&\quad - (2g(I - T_{hp})^{-1}f - fg) \\
&\quad + 2g(I - T_{hp})^{-1}g - g^2 \\
&= (I - T_{h^2p})^{-1}(2(f - g)(I - T_{hp})^{-1}(f - g) - (f - g)^2).
\end{aligned} \tag{12}$$

It is convenient to rewrite w_1 as

$$w_1 = w_{11} + w_{12}$$

with

$$w_{11} = ((I - T_{k^2p})^{-1} - 2(I - T_{khp})^{-1} + (I - T_{h^2p})^{-1})(2f(I - T_{kp})^{-1}f - f^2) \tag{13}$$

and

$$\begin{aligned}
w_{12} &= (I - T_{khp})^{-1}(2f(I - T_{kp})^{-1}f - 2f(I - T_{hp})^{-1}f) \\
&\quad + (I - T_{h^2p})^{-1}(2f(I - T_{hp})^{-1}f - 2f(I - T_{kp})^{-1}f) \\
&= ((I - T_{khp})^{-1} - (I - T_{h^2p})^{-1})(2f((I - T_{kp})^{-1} - (I - T_{hp})^{-1})f) \\
&= (I - T_{khp})^{-1}(T_{khp} - T_{h^2p})(I - T_{h^2p})^{-1} \\
&\quad (2f(I - T_{kp})^{-1}(T_{kp} - T_{hp})(I - T_{hp})^{-1}f) \\
&= (I - T_{khp})^{-1}T_{(k-h)hp}(I - T_{h^2p})^{-1} \\
&\quad (2f(I - T_{kp})^{-1}T_{(k-h)p}(I - T_{hp})^{-1}f).
\end{aligned} \tag{14}$$

We need relation (27) of [Hei95]:

$$\|T_{(k-h)hp} : L_\infty(X) \rightarrow L_\infty(X)\| = \|(k-h)hp\|_{L_\infty(L_1)} \leq \alpha\delta(k, h).$$

With the notation of Lemma 1 we get from (12) and (14)

$$\begin{aligned} \|w_2\|_{L_\infty(X)} &\leq \beta_0(2\beta_1 + 1)\|f - g\|_{L_\infty(X)}^2 \\ \|w_{12}\|_{L_\infty(X)} &\leq \beta_0\alpha\delta(k, h)\beta_02\theta\beta_1\delta(k, h)\beta_1\theta \\ &= 2\alpha\beta_0^2\beta_1^2\theta^2\delta(k, h)^2 \leq 2\alpha\beta_0^2\beta_1^2\theta^2\varepsilon(k, h)^2. \end{aligned}$$

For the estimate of (13) we recall relations (26) and (28) from [Hei95]

$$\begin{aligned} &(I - T_{k^2p})^{-1} - 2(I - T_{khp})^{-1} + (I - T_{h^2p})^{-1} \\ &= (I - T_{k^2p})^{-1}(T_{(k-h)^2p} + T_{(h^2-k^2)p}(I - T_{h^2p})^{-1}T_{(h-k)hp})(I - T_{khp})^{-1} \end{aligned} \quad (15)$$

and

$$\|T_{(h^2-k^2)p} : L_\infty(X) \rightarrow L_\infty(X)\| \leq 2\alpha\delta(k, h).$$

This gives

$$\begin{aligned} \|w_{11}\| &\leq \beta_0 \left(\varepsilon(k, h)^2 + 2\alpha^2\beta_0\delta(k, h)^2 \right) \beta_0 \|2f(I - T_{kp})^{-1}f - f^2\|_{L_\infty(X)} \\ &\leq \beta_0^2 \left(\varepsilon(k, h)^2 + 2\alpha^2\beta_0\delta(k, h)^2 \right) (1 + 2\beta_1)\theta^2 \\ &\leq \beta_0^2\theta^2(1 + 2\alpha^2\beta_0)(1 + 2\beta_1)\varepsilon(k, h)^2. \end{aligned}$$

This completes the proof.

Finally, we show that in analogy to Theorem 5 of [Hei95], L_2 -estimates can be obtained for the variance, once slightly stronger assumptions are imposed. We suppose that $\varphi^2p_0 \in L_\infty(X)$ and

$$\operatorname{ess\,sup}_{y \in X} \int_X p(x, y) \, d\mu(x) \leq 1 \quad (16)$$

which is the case, in particular, if $p(x, y)$ is symmetric (as e.g. the choice of the transition probability for the radiance equation in [Hei95]). Define for $\alpha > 0$, $0 < \gamma, \gamma_1 < 1$, $n_0, n_1 \in \mathbf{N}$

$$\mathcal{K}^*(\alpha, \gamma, \gamma_1, n_0, n_1) = \left\{ k \in \mathcal{K}(\alpha, \gamma, n_0) : \|T_{k^2p}^{n_1} : L_1(X) \rightarrow L_1(X)\| \leq \gamma_1 \right\}.$$

Lemma 4 Let $\alpha > 0$, $0 < \gamma, \gamma_1 < 1$, $n_0, n_1 \in \mathbf{N}$. If $k, h \in \mathcal{K}(\alpha, \gamma, \gamma_1, n_0, n_1)$, then

(i) $I - T_{k^2p}$ is invertible in $L_1(X)$ and $\|(I - T_{k^2p})^{-1} : L_1(X) \rightarrow L_1(X)\| \leq \beta_2$ with $\beta_2 = (1 - \gamma_1)^{-1} \sum_{j=0}^{n_1-1} \alpha^{2j}$,

(ii) $I - T_{kp}$ is invertible in $L_1(X)$ and $\|(I - T_{kp})^{-1} : L_1(X) \rightarrow L_1(X)\| \leq \beta_3$, with $\beta_3 = (1 - \gamma_1)^{-1} \sum_{j=0}^{n_1-1} \alpha^j$,

(iii) $I - T_{khp}$ is invertible in $L_1(X)$ and $\|(I - T_{khp})^{-1} : L_1(X) \rightarrow L_1(X)\| \leq \beta_2$.

Proof: For an integral operator T_K we have

$$\|T_K : L_1(X) \rightarrow L_1(X)\| = \|T_{K^*} : L_\infty(X) \rightarrow L_\infty(X)\|,$$

where $K^*(x, y) = K(y, x)$. Hence, if $k, h \in \mathcal{K}(\alpha, \gamma, \gamma_1, n_0, n_1)$, then $k^*, h^* \in \mathcal{K}_{p^*}(\alpha, \gamma_1, n_1)$ where \mathcal{K}_{p^*} denotes the class \mathcal{K} , based on p^* instead of p . Now Lemma 4 is a direct consequence of Lemma 1.

Finally, we denote

$$\sigma(k, h)^2 = \int_X \int_X (k(x, y) - h(x, y))^2 p(x, y) d\mu(x) d\mu(y).$$

Theorem 5 Given $\alpha, \theta > 0$, $0 < \gamma, \gamma_1 < 1$, $n_0, n_1 \in \mathbf{N}$, there exists a constant $c > 0$ such that for $k, h \in \mathcal{K}^*(\alpha, \gamma, \gamma_1, n_0, n_1)$ and $f, g \in \mathcal{F}(\theta)$,

$$\text{Var}(\zeta(k, f, h, g, \xi)) \leq c \left(\sigma(k, h)^2 + \|f - g\|_{L_2(X)}^2 \right).$$

Proof: Lemma 1 and 4 yield

$$\begin{aligned} \|(I - T_{h^2p})^{-1} : L_\infty(X) \rightarrow L_\infty(X)\| &\leq \beta_0, \text{ and} \\ \|(I - T_{h^2p})^{-1} : L_1(X) \rightarrow L_1(X)\| &\leq \beta_2 \end{aligned}$$

and hence, by the Riesz-Thorin interpolation theorem (see [Tri78]),

$$\|(I - T_{h^2p})^{-1} : L_2(X) \rightarrow L_2(X)\| \leq (\beta_0 \beta_2)^{1/2}. \quad (17)$$

Using (10 – 14) of the previous proof, we obtain

$$\text{Var}(\zeta) \leq 2 \left(\|w_{11}\|_{L_1(X)} + \|w_{12}\|_{L_1(X)} + \|w_2\|_{L_1(X)} \right) \|\varphi^2 p_0\|_{L_\infty(X)}.$$

First we estimate $\|w_{11}\|_{L_1(X)}$ on the basis of relations (13) and (15). From (17) of the present paper and inequalities (35 – 37) of [Hei95] we get

$$\begin{aligned} \|T_{(k-h)^2p} : L_\infty(X) \rightarrow L_1(X)\| &\leq \sigma(k, h)^2 \\ \|T_{(h^2-k^2)p}(I - T_{h^2p})^{-1}T_{(h-k)hp} : L_\infty(X) \rightarrow L_1(X)\| &\leq 2\alpha^2(\beta_0\beta_2)^{1/2}\sigma(k, h)^2. \end{aligned}$$

This gives

$$\begin{aligned} \|(I - T_{k^2p})^{-1} - 2(I - T_{khp})^{-1} + (I - T_{h^2p})^{-1} : L_\infty(X) \rightarrow L_1(X)\| \\ \leq \beta_0\beta_2(2\alpha^2(\beta_0\beta_2)^{1/2} + 1)\sigma(k, h)^2. \end{aligned}$$

Moreover, from Lemma 1,

$$\|2f(I - T_{kp})^{-1}f - f^2\|_{L_\infty(X)} \leq (2\beta_1 + 1)\theta^2.$$

The last two relations combined with (13) yield

$$\|w_{11}\|_{L_1(X)} \leq \beta_0\beta_2\theta^2(2\beta_1 + 1)(2\alpha^2(\beta_0\beta_2)^{1/2} + 1)\sigma(k, h)^2.$$

Next we deal with w_{12} . From (14) we infer

$$\begin{aligned} \|w_{12}\|_{L_1(X)} &\leq \|(I - T_{khp})^{-1}T_{(k-h)hp}(I - T_{h^2p})^{-1} : L_2(X) \rightarrow L_1(X)\| \\ &\quad \times \|2f(I - T_{kp})^{-1}T_{(k-h)p}(I - T_{hp})^{-1}f\|_{L_2(X)}. \end{aligned} \tag{18}$$

We shall use (36) and (37) of [Hei95] which give

$$\|T_{(k-h)hp} : L_2(X) \rightarrow L_1(X)\| \leq \alpha\sigma(k, h).$$

Consequently, the first factor of (18) satisfies

$$\begin{aligned}
& \|(I - T_{khp})^{-1}T_{(k-h)hp}(I - T_{h^2p})^{-1} : L_2(X) \rightarrow L_1(X)\| \\
& \leq \|(I - T_{khp})^{-1} : L_1(X) \rightarrow L_1(X)\| \cdot \|T_{(k-h)hp} : L_2(X) \rightarrow L_1(X)\| \\
& \quad \times \|(I - T_{h^2p})^{-1} : L_2(X) \rightarrow L_2(X)\| \\
& \leq \alpha\beta_0^{1/2}\beta_2^{3/2}\sigma(k, h).
\end{aligned}$$

To estimate the second factor of (18), we note that (36) of [Hei95] gives

$$\|T_{(k-h)p} : L_\infty(X) \rightarrow L_2(X)\| \leq \sigma(k, h).$$

Moreover, arguing in the same way as for the derivation of (17), we conclude from Lemmas 1 and 4

$$\|(I - T_{kp})^{-1} : L_2(X) \rightarrow L_2(X)\| \leq (\beta_1\beta_3)^{1/2}.$$

Hence,

$$\begin{aligned}
& \|2f(I - T_{kp})^{-1}T_{(k-h)p}(I - T_{hp})^{-1}f\|_{L_2(X)} \\
& \leq 2\theta^2\|(I - T_{kp})^{-1}T_{(k-h)p}(I - T_{hp})^{-1} : L_\infty(X) \rightarrow L_2(X)\| \\
& \leq 2\theta^2\|(I - T_{kp})^{-1} : L_2(X) \rightarrow L_2(X)\| \|T_{(k-h)p} : L_\infty(X) \rightarrow L_2(X)\| \\
& \quad \times \|(I - T_{hp})^{-1} : L_\infty(X) \rightarrow L_\infty(X)\| \\
& \leq 2\theta^2\beta_1^{3/2}\beta_3^{1/2}\sigma(k, h).
\end{aligned}$$

So we get

$$\|w_{12}\|_{L_1(X)} \leq 2\alpha\beta_0^{1/2}\beta_1^{3/2}\beta_2^{3/2}\beta_3^{1/2}\theta^2\sigma(k, h)^2.$$

Finally, we turn to w_2 . According to (12),

$$\begin{aligned}
\|w_2\|_{L_1(X)} & \leq \|(I - T_{h^2p})^{-1} : L_1(X) \rightarrow L_1(X)\| \\
& \quad \times \|2(f - g)(I - T_{hp})^{-1}(f - g) - (f - g)^2\|_{L_1(X)} \\
& \leq \beta_2 \left(2\|(I - T_{hp})^{-1} : L_2(X) \rightarrow L_2(X)\| \|f - g\|_{L_2(X)}^2 + \|f - g\|_{L_2(X)}^2 \right) \\
& \leq \beta_2(2(\beta_1\beta_3)^{1/2} + 1)\|f - g\|_{L_2(X)}^2.
\end{aligned}$$

This completes the proof.

In conclusion, let us mention a few consequences of the results proved above. They are of the same form (up to obvious modifications) as those in section 3 of [Hei95]. It follows that all consequences drawn in [Hei95] hold true also for the collision estimate. In particular, Corollary 4 is valid, which shows that the optimal rate obtained in [HM93] for the absorption estimate is also true for the collision estimate. Moreover, the analysis of section 4 of [Hei95] carries over: Using some approximate deterministic Galerkin solution, one can construct neighboring h and g of increasing precision, resulting in decreasing variance of the new scheme based on the collision estimate. To estimate the variance, one can use Propositions 6 and 7 of [Hei95]. Finally, the applications to the radiance equation of computer graphics hold true as well. Details can be found in [Hei95] and [Hei96]

References

- [EM82] S. M. Ermakov and G. A. Mikhailov. *The Theory of Statistical Trials* (in Russian). Nauka, Moscow, 1982.
- [ENS89] S. M. Ermakov, W. W. Nekrutkin, and A. S. Sipin. *Random Processes for Classical Equations of Mathematical Physics*, volume 34 of Math. Appl. (Soviet Ser.). Kluwer, Dodrecht, 1989.
- [Erm71] S. M. Ermakov. *The Monte Carlo Method and Related Problems* (in Russian). Nauka, Moscow, 1971.
- [ES85] S. M. Ermakov and A. S. Sipin. A new scheme of the Monte Carlo method for solving problems of mathematical physics. *Soviet Phys. Doklady*, **30**(11): 935 – 936, 1985.
- [Hei94] S. Heinrich. Random approximation in numerical analysis. In K. D. Bierstedt, A. Pietsch, W. M. Ruess, and D. Vogt, editors, *Functional Analysis*, pages 123 – 171. Marcel Dekker, 1994.
- [Hei95] S. Heinrich. Variance reduction for Monte Carlo methods by means of deterministic numerical computation. *Monte Carlo Methods and Appl.*, **1**: 251 – 277, 1995.
- [Hei96] S. Heinrich. Using the Galerkin method for variance reduction. In *2nd. St. Petersburg Workshop on Simulation*, pages 56 – 59. Publishing House of St. Petersburg University, 1996.
- [HM93] S. Heinrich and P. Mathé. The Monte Carlo complexity of Fredholm integral equations. *Math. of Computation*, **60**: 257 – 278, 1993.
- [KW86] M. H. Kalos and P. A. Whitlock. *Monte Carlo Methods, Vol. I: Basics*. Wiley, New York, 1986.

- [LW94] E. P. Lafortune and Y. D. Willems. The ambient term as a variance reducing technique for Monte Carlo ray tracing. In *Proc. 5th Eurographics Workshop*. Darmstadt, 1994.
- [Mik91a] G. A. Mikhailov. *Minimization of Computational Costs of Non-Analogue Monte Carlo Methods*. World Scientific, Singapore, 1991.
- [Mik91b] G. A. Mikhailov. *Optimization of Weighted Monte Carlo Methods*. Springer, 1991.
- [Sab91] K. K. Sabelfeld. *Monte Carlo Methods in Boundary Value Problems*. Springer, Berlin-Heidelberg-New York, 1991.
- [SG69] J. Spanier and E. M. Gelbard. *Monte Carlo Principles and Neutron Transport Problems*. Addison-Wesley, Reading, Massachusetts, 1969.
- [Sob73] I. M. Sobol. *Computational Monte Carlo Methods* (in Russian). Nauka, Moscow, 1973.
- [Spa79] J. Spanier. A new family of estimators for random walk problems. *J. Inst. Math. Appl.*, **23**: 1 – 31, 1979.
- [Tri78] H. Triebel. *Interpolation Theory, Function Spaces, Differential Operators*. Deutscher Verlag der Wissenschaften, Berlin, 1978.