



Fraunhofer Institut
Techno- und
Wirtschaftsmathematik

H. Knaf

Kernel Fisher discriminant functions –
a concise and rigorous introduction

© Fraunhofer-Institut für Techno- und Wirtschaftsmathematik ITWM 2007

ISSN 1434-9973

Bericht 117 (2007)

Alle Rechte vorbehalten. Ohne ausdrückliche schriftliche Genehmigung des Herausgebers ist es nicht gestattet, das Buch oder Teile daraus in irgendeiner Form durch Fotokopie, Mikrofilm oder andere Verfahren zu reproduzieren oder in eine für Maschinen, insbesondere Datenverarbeitungsanlagen, verwendbare Sprache zu übertragen. Dasselbe gilt für das Recht der öffentlichen Wiedergabe.

Warennamen werden ohne Gewährleistung der freien Verwendbarkeit benutzt.

Die Veröffentlichungen in der Berichtsreihe des Fraunhofer ITWM können bezogen werden über:

Fraunhofer-Institut für Techno- und
Wirtschaftsmathematik ITWM
Fraunhofer-Platz 1

67663 Kaiserslautern
Germany

Telefon: +49 (0) 6 31/3 16 00-0
Telefax: +49 (0) 6 31/3 16 00-1099
E-Mail: info@itwm.fraunhofer.de
Internet: www.itwm.fraunhofer.de

Vorwort

Das Tätigkeitsfeld des Fraunhofer-Instituts für Techno- und Wirtschaftsmathematik ITWM umfasst anwendungsnahe Grundlagenforschung, angewandte Forschung sowie Beratung und kundenspezifische Lösungen auf allen Gebieten, die für Techno- und Wirtschaftsmathematik bedeutsam sind.

In der Reihe »Berichte des Fraunhofer ITWM« soll die Arbeit des Instituts kontinuierlich einer interessierten Öffentlichkeit in Industrie, Wirtschaft und Wissenschaft vorgestellt werden. Durch die enge Verzahnung mit dem Fachbereich Mathematik der Universität Kaiserslautern sowie durch zahlreiche Kooperationen mit internationalen Institutionen und Hochschulen in den Bereichen Ausbildung und Forschung ist ein großes Potenzial für Forschungsberichte vorhanden. In die Berichtreihe sollen sowohl hervorragende Diplom- und Projektarbeiten und Dissertationen als auch Forschungsberichte der Institutsmitarbeiter und Institutsgäste zu aktuellen Fragen der Techno- und Wirtschaftsmathematik aufgenommen werden.

Darüber hinaus bietet die Reihe ein Forum für die Berichterstattung über die zahlreichen Kooperationsprojekte des Instituts mit Partnern aus Industrie und Wirtschaft.

Berichterstattung heißt hier Dokumentation des Transfers aktueller Ergebnisse aus mathematischer Forschungs- und Entwicklungsarbeit in industrielle Anwendungen und Softwareprodukte – und umgekehrt, denn Probleme der Praxis generieren neue interessante mathematische Fragestellungen.



Prof. Dr. Dieter Prätzel-Wolters
Institutsleiter

Kaiserslautern, im Juni 2001

Kernel Fisher discriminant functions – a concise and rigorous introduction

Hagen Knaf

September 2006

Abstract

In the article the application of kernel functions – the so-called »kernel trick« – in the context of Fisher’s approach to linear discriminant analysis is described for data sets subdivided into two groups and having real attributes. The relevant facts about functional Hilbert spaces and kernel functions including their proofs are presented. The approximative algorithm published in [Mik3] to compute a discriminant function given the data and a kernel function is briefly reviewed. As an illustration of the technique an artificial data set is analysed using the algorithm just mentioned.

Keywords: discriminant analysis, functional Hilbert space, kernel function, reproducing kernel.

MSC: 46E22, 46N30, 62-07.

Introduction

A fundamental problem in data mining consists of separating two disjoint, finite subsets X_{-1} and X_{+1} of the euclidean space \mathbb{R}^m through a hypersurface $S \subset \mathbb{R}^m$. More precisely and in analytic terms this amounts to choosing a function

$$d_{\theta^*} : \mathbb{R}^m \rightarrow \mathbb{R}$$

within a given parametrized family

$$F = \{d_{\theta} : \mathbb{R}^m \rightarrow \mathbb{R} \mid \theta \in \Theta\}, \quad \Theta \subseteq \mathbb{R}^p,$$

of functionals, such that

$$X_k \subseteq S_k, \quad k \in \{-1, +1\}, \quad (1)$$

where $S_k := \{x \in \mathbb{R}^m \mid \text{sign}(d_{\theta^*}(x)) = k\}$ are the half-spaces defined by $S := d_{\theta^*}^{-1}(0)$. Of course depending on F such a function d_{θ^*} needs not exist. One therefore replaces the requirement (1) by the weaker

$$\text{maximize } \text{obj}(X_{-1} \cap S_{-1}, X_{+1} \cap S_{+1}), \quad (2)$$

where obj is an objective function like for example the »total hitrate«

$$\text{obj}(X_{-1} \cap S_{-1}, X_{+1} \cap S_{+1}) := \frac{|X_{-1} \cap S_{-1}| + |X_{+1} \cap S_{+1}|}{|X_{-1}| + |X_{+1}|}.$$

In [Fis] Fisher showed that within the family

$$F = \{d_{(a,c)}(x) := \langle x, a \rangle + c \mid a \in \mathbb{R}^m, c \in \mathbb{R}\} \quad (3)$$

there exists a function $d_{(a^*, c^*)}$ that maximizes the objective function

$$\text{obj}(a, c) := \frac{\|p(\bar{x}_{-1}) - p(\bar{x}_{+1})\|^2}{\bar{s}_{-1}^2 + \bar{s}_{+1}^2}, \quad (4)$$

where $p(\bar{x}_i)$ and \bar{s}_i^2 are the mean values and variances of the sets X_{-1} and X_{+1} after orthogonal projection to a line perpendicular to the hyperplane $d_{(a,c)}^{-1}(0)$.

The optimal parameters (a^*, c^*) can be determined analytically and are unique if the set $X := X_{-1} \cup X_{+1}$ is in a »sufficiently general« position.

Only some years ago it was realized that the particular form of Fisher's objective function (4) allows to apply the so-called »kernel trick« to obtain a non-linear version of Fisher's discriminant function $d_{(a^*, c^*)}$: instead of (3) one considers families of the form

$$F = \left\{ \sum_{y \in X} a_y K(x, y) + c \mid a_y, c \in \mathbb{R} \right\},$$

where $K : \mathbb{R}^m \times \mathbb{R}^m \rightarrow \mathbb{R}$ is a so-called kernel function. The particular properties of K allow to construct a Hilbert space H consisting of functions $h : X \rightarrow \mathbb{R}$ such that H is generated by the elements of $Z := \{K(\cdot, y) \mid y \in X\}$. Solving the optimization problem (2) using Fisher's objective function (4) in the space H and for the data set $Z = Z_{-1} \cup Z_{+1}$, where $Z_k := \{K(\cdot, y) \mid y \in X_k\}$, yields a in general nonlinear discriminant function

$$d^*(x) = \sum_{y \in X} a_y^* K(x, y) + c^*.$$

In the present article a concise and (hopefully) rigorous presentation of this whole process is provided, taking an analytic point of view.

1 Discriminant Analysis

In this section a brief review of the general problem of discriminant analysis is given, followed by a summary of relevant facts about Fisher's linear discriminant function.

1.1 The problem

Let Ω be a set that is decomposed into $g \geq 2$ pairwise disjoint subsets Ω_i , in the sequel called **groups**:

$$\Omega = \Omega_1 \cup \dots \cup \Omega_g. \quad (5)$$

This decomposition yields the **label map**

$$\ell : \Omega \rightarrow \{1, \dots, g\}, (\omega \in \Omega_k) \mapsto k, \quad (6)$$

that assigns to an object $\omega \in \Omega$ its group label.

Every object $\omega \in \Omega$ comes equipped with a finite set of **attributes** $x(\omega)$. To keep the overview focussed on the facts relevant for the present article it is assumed that these attributes are real numbers, thus giving the **attribute map**

$$\Omega \rightarrow A \subseteq \mathbb{R}^m, \omega \mapsto x(\omega). \quad (7)$$

Based on the known attributes and group labels of a set of samples $\Lambda \subset \Omega$ randomly drawn from Ω in discriminant analysis one seeks to determine an estimate

$$\hat{\ell} : \Omega \rightarrow \{1, \dots, g\} \quad (8)$$

of the unknown label function ℓ . Since by assumption the estimated label $\hat{\ell}(\omega)$ depends on $x(\omega)$ only, determining $\hat{\ell}$ amounts to determine a **decision function**

$$\delta : A \rightarrow \{1, \dots, g\} \quad (9)$$

such that $\hat{\ell}(\omega) = \delta(x(\omega))$.

A typical example is the medical diagnosis of a specific disease based on a blood test say. Here the set Ω consists of a population of human beings, the attributes $x(\omega)$ are the values of the parameters measured during the blood test and the label $\ell(\omega)$ encodes whether the human ω is suffering of the considered disease ($\ell(\omega) = 1$ say) or not ($\ell(\omega) = 2$). The estimated label $\hat{\ell}(\omega)$ represents the diagnosis based on the blood test. It can be correct or false. Note that in this situation the map $\omega \mapsto x(\omega)$ needs not be injective, so that the diagnostic performance is limited already by the choice of the particular parameters measured during the blood test.

From now on only the case $g = 2$ is considered. Problems with $g > 2$ can be treated by successively splitting off one group after the other. Using the algorithm presented in Section 3 this can be done effectively.

For reasons that will become clear in the next paragraph it is assumed that the group labels for the two groups are -1 and $+1$. Moreover a decision function is allowed to take the value 0 with the meaning that for an object $\omega \in \Omega$ with the property $\delta(x(\omega)) = 0$ the actual group label $\ell(\omega)$ cannot be estimated using the decision function δ .

An intuitive geometric way to determine a (modified) decision function

$$\delta : A \rightarrow \{-1, 0, 1\} \quad (10)$$

is to fix a hypersurface $S \subset \mathbb{R}^m$ such that most of the samples in the set $X_{-1} := \{x(\omega) \mid \omega \in \Lambda \cap \Omega_{-1}\}$ »lie on one side of S «, while most of the samples in the set $X_{+1} := \{x(\omega) \mid \omega \in \Lambda \cap \Omega_{+1}\}$ »lie on the other side«. This approach can be made more precise in the following way: a function $d : \mathbb{R}^m \rightarrow \mathbb{R}$ yields a hypersurface through

$$S := d^{-1}(0)$$

provided it for example satisfies the prerequisites of the Implicit Function Theorem. The decision function associated to that hypersurface in the sense vaguely described above is then given by

$$\delta : A \rightarrow \{-1, 0, 1\}, \quad x \mapsto \text{sign}(d(x)), \quad (11)$$

where sign denotes the function that maps a real number to its sign. The function d is called a **discriminant function for the grouping** $X_{-1} \cup X_{+1}$.

A strategy to choose a discriminant function is to consider a sufficiently general family

$$F := \{d_\theta : \mathbb{R}^m \rightarrow \mathbb{R} \mid \theta \in \Theta\}, \quad \Theta \subseteq \mathbb{R}^p, \quad (12)$$

of functionals possessing the necessary properties to define hypersurfaces via $S := d_\theta^{-1}(0)$ and to choose a parameter θ^* optimal with respect to some objective function obj derived from the known restriction $\ell|_\Lambda$ of the label map.

1.2 Fisher's linear discriminant function

Let Λ be a finite set of samples drawn from the population $\Omega = \Omega_{-1} \cup \Omega_{+1}$ grouped into two groups with labels -1 and $+1$. The attribute map then yields

$$x(\Lambda) =: X = X_{-1} \cup X_{+1} \subset \mathbb{R}^m, \quad X_k := x(\Lambda \cap \Omega_k), \quad k \in \{-1, +1\}. \quad (13)$$

From now on and throughout the whole article it is assumed that the attribute map is injective. Thus (13) is a subdivision of X into two disjoint groups.

Among the various ways to choose an **affine discriminant function**

$$d(x) = f(x) + c, \quad f \in \text{Hom}(\mathbb{R}^m, \mathbb{R}), \quad c \in \mathbb{R}, \quad (14)$$

to separate the two groups in (13) R. A. Fisher's approach is known to be robust and has the advantage of leading to an analytic solution: for a one-dimensional sub-vector space $L \subseteq \mathbb{R}^m$ consider the orthogonal projection $p_L : \mathbb{R}^m \rightarrow L$ and define the **Fisher discriminant of $X_{-1} \cup X_{+1}$ with respect to L** as:

$$\mathcal{F}(L) := \frac{\|p_L(\bar{x}_{-1}) - p_L(\bar{x}_{+1})\|^2}{\bar{s}_{-1}^2 + \bar{s}_{+1}^2}, \quad (15)$$

where the

$$\bar{x}_k := \frac{1}{|X_k|} \sum_{x \in X_k} x$$

are the **group centroids** and

$$\bar{s}_k^2 := \sum_{x \in X_k} \|p_L(x) - p_L(\bar{x}_k)\|^2, \quad (16)$$

up to a factor are the variances of the sets $p_L(X_k)$. Fisher proposes to consider the discriminant function

$$d^*(x) = p_{L^*}(x) + c^*, \quad (17)$$

where L^* is a solution of the optimization problem

$$\max(\mathcal{F}(L) \mid L \subseteq \mathbb{R}^m \text{ a one-dimensional sub-vector space}) \quad (18)$$

and

$$c^* := -\frac{1}{2}(p_{L^*}(\bar{x}_{-1}) + p_{L^*}(\bar{x}_{+1})). \quad (19)$$

Note that $-c^*$ is the mean value of the centroids of the projected groups $p_{L^*}(X_k)$.

In Section 3 a non-linear variant of Fisher's discriminant function is constructed using the so-called »kernel trick«. This is possible because a solution L^* of the optimization problem (18) can be obtained from quantities, that can be expressed solely in terms of scalar products between the elements $x \in X$. To verify this property one has to make the reasonable assumption

$$\sum_{x \in X} \mathbb{R}x = \mathbb{R}^m. \quad (20)$$

The orthogonal projection p_L to the one-dimensional sub-vector space $L \subseteq \mathbb{R}^m$ can be expressed as $p_L(x) = \langle a, x \rangle a$ with some $a \in \mathbb{R}^m$ having the property $\|a\| = 1$. Using (20) one can write

$$a = \sum_{j=1}^n a_j x_j, \quad a_j \in \mathbb{R},$$

where $\{x_1, \dots, x_n\} = X$ is some numbering of the elements of X . Defining the vectors

$$\alpha := (a_1, \dots, a_n) \in \mathbb{R}^n$$

and

$$m_k := (\langle x_1, \bar{x}_k \rangle, \dots, \langle x_n, \bar{x}_k \rangle), \quad k \in \{-1, +1\}, \quad (21)$$

one gets

$$\begin{aligned} \|p_L(\bar{x}_{-1}) - p_L(\bar{x}_{+1})\|^2 &= \left\langle \sum_{j=1}^n a_j x_j, \bar{x}_{-1} - \bar{x}_{+1} \right\rangle^2 \\ &= \left(\sum_{j=1}^n a_j \langle x_j, \bar{x}_{-1} - \bar{x}_{+1} \rangle \right)^2 \\ &= (\alpha^t (m_{-1} - m_{+1}))^2 \end{aligned}$$

and thus the relation

$$\|p_L(\bar{x}_{-1}) - p_L(\bar{x}_{+1})\|^2 = \alpha^t (m_{-1} - m_{+1})(m_{-1} - m_{+1})^t \alpha \quad (22)$$

for the numerator of (15). The entries of the matrix

$$B := (m_{-1} - m_{+1})(m_{-1} - m_{+1})^t \in \mathbb{R}^{n \times n} \quad (23)$$

are sums of products of scalar products between elements of X . The denominator of (15) can be written in a similar way:

$$\begin{aligned} \bar{s}_{-1}^2 + \bar{s}_{+1}^2 &= \sum_{k \in \{-1, +1\}} \sum_{x \in X_k} \langle a, x \rangle^2 + \langle a, \bar{x}_k \rangle^2 - 2\langle a, x \rangle \langle a, \bar{x}_k \rangle \\ &= \left(\sum_{j=1}^n \langle a, x_j \rangle^2 \right) - |X_{-1}| \langle a, \bar{x}_{-1} \rangle^2 - |X_{+1}| \langle a, \bar{x}_{+1} \rangle^2 \\ &= \left(\sum_{j=1}^n \langle a, x_j \rangle^2 \right) - |X_{-1}| \alpha^t m_{-1} m_{-1}^t \alpha - |X_{+1}| \alpha^t m_{+1} m_{+1}^t \alpha. \end{aligned}$$

Using the matrix

$$K := (\langle x_i, x_j \rangle)_{i,j \in \{1, \dots, n\}} \in \mathbb{R}^{n \times n} \quad (24)$$

the first summand on the right hand side can also be written as a matrix product:

$$\sum_{j=1}^n \langle a, x_j \rangle^2 = \alpha^t K K^t \alpha.$$

The optimization problem (18) can now be reformulated: the solutions L^* of (18) correspond to the solutions $\alpha^* = (a_1^*, \dots, a_n^*)$ of the optimization problem

$$\max \left(\frac{\alpha^t B \alpha}{\alpha^t W \alpha} \mid \alpha \in \mathbb{R}^n \setminus 0 \right) \quad (25)$$

through setting $p_{L^*}(x) := \langle a^*, x \rangle a^*$, $a^* := \sum_{j=1}^n a_j^* x_j$. Here the matrix W is defined as:

$$W := K K^t - |X_{-1}| m_{-1} m_{-1}^t - |X_{+1}| m_{+1} m_{+1}^t \in \mathbb{R}^{n \times n}. \quad (26)$$

It is well-known that (25) possesses the unique solution

$$\alpha^* := W^{-1}(m_{-1} - m_{+1}) \quad (27)$$

provided that W is invertible.

The constant term c^* can be expressed in terms of scalar products between the elements of X and the solution α^* :

$$c^* = -\frac{1}{2} (\alpha^*)^t (m_{-1} + m_{+1}). \quad (28)$$

2 Hilbert spaces from kernel functions

In this section an exposition of those parts of the theory of functional Hilbert spaces is given, that lead to the following statement sometimes called the »kernel trick«:

every real- or complex-valued function $K : X \times X \rightarrow \mathbb{K}$ with the properties

- $\forall x_1, x_2 \in X : K(x_1, x_2) = \overline{K(x_2, x_1)}$,
- $\forall n \in \mathbb{N}, \forall (x_1, \dots, x_n) \in X^n : (K(x_i, x_j))_{i,j \in \{1, \dots, n\}} \in \mathbb{K}^{n \times n}$ is positive semi-definite,

gives rise to a Hilbert space $H(K)$ consisting of functions $h : X \rightarrow \mathbb{K}$, and to a map $\phi : X \rightarrow H(K)$ such that

$$\forall x_1, x_2 \in X : \langle \phi(x_1), \phi(x_2) \rangle = K(x_2, x_1),$$

where $\langle \cdot, \cdot \rangle$ denotes the scalar product of $H(K)$.

2.1 Functional Hilbert spaces

For a set X the set $\text{Map}(X, \mathbb{K})$ of all maps $h : X \rightarrow \mathbb{K}$ into a field \mathbb{K} becomes a vector space over \mathbb{K} if addition and scalar multiplication are defined pointwise. The maps $h_x \in \text{Map}(X, \mathbb{K})$ defined by $h_x(x) = 1$ and $h_x(y) = 0$ for $x \neq y$ are linearly independent and form a basis of $\text{Map}(X, \mathbb{K})$ if X is finite. Hence $\text{Map}(X, \mathbb{K})$ has finite dimension if and only if X is finite.

Every $x \in X$ gives rise to the \mathbb{K} -linear **evaluation functional**

$$e_x : \text{Map}(X, \mathbb{K}) \rightarrow \mathbb{K}, h \mapsto h(x). \quad (29)$$

These maps play a particular role in the sequel.

From now on let \mathbb{K} be either the field of reals or the field of complex numbers.

A Hilbert space H over \mathbb{K} is called **functional over X** if it is a sub-vector space of $\text{Map}(X, \mathbb{K})$. If H has infinite dimension, the evaluation functionals (29) may not be continuous. However from now on only functional Hilbert spaces with the property that *all* evaluation functionals are continuous are considered. In particular norm convergence in H implies pointwise convergence:

$$\left(\lim_{k \rightarrow \infty} h_k = h \right) \Rightarrow \left(\forall x \in X : \lim_{k \rightarrow \infty} h_k(x) = h(x) \right). \quad (30)$$

The principal result of the structure theory of Hilbert spaces states that every Hilbert space H is isomorphic to a space $\ell^2(X)$ of square-summable maps

$X \rightarrow \mathbb{K}$. In particular it follows that every Hilbert space is functional over some set X , even with continuous evaluation functionals.

Proposition 2.1 *For a Hilbert space H functional over X and possessing continuous evaluation functionals there exists a unique function $K : X \times X \rightarrow \mathbb{K}$ with the properties:*

$$\begin{aligned} \forall x \in X : \quad K_x &:= K(\cdot, x) \in H, \\ e_x &= \langle \cdot, K_x \rangle. \end{aligned} \tag{31}$$

The function K is called the **reproducing kernel of H** . It has the following properties:

- (1) $\forall x_1, x_2 \in X : K(x_1, x_2) = \overline{K(x_2, x_1)}$,
- (2) $\forall n \in \mathbb{N}, \forall (x_1, \dots, x_n) \in X^n : K(x_1, \dots, x_n) := (K(x_i, x_j))_{i,j \in \{1, \dots, n\}}$ is positive semi-definite, that is

$$\forall z \in \mathbb{K}^n : z^t K(x_1, \dots, x_n) \bar{z} \geq 0.$$

Proof. By the Representation Theorem of Riesz for every $x \in X$ there exists a unique element $K_x \in H$ such that $e_x = \langle \cdot, K_x \rangle$. Thus one can define the map K through

$$K(x, y) := K_y(x). \tag{32}$$

The uniqueness of K now also follows.

By definition K satisfies

$$K(x, y) = K_y(x) = \langle K_y, K_x \rangle, \tag{33}$$

which implies the symmetry of K . As for positive semi-definiteness this relation yields

$$\begin{aligned} z^t K(x_1, \dots, x_n) \bar{z} &= \sum_{j=1}^n \sum_{k=1}^n z_j K(x_j, x_k) \bar{z}_k \\ &= \left\langle \sum_{k=1}^n \bar{z}_k K_{x_k}, \sum_{j=1}^n z_j K_{x_j} \right\rangle \geq 0, \end{aligned}$$

where $z^t = (z_1, \dots, z_n) \in \mathbb{K}^n$. □

As a consequence of Proposition 2.1 for a Hilbert space H functional over X and having continuous evaluation functionals one can define a map

$$\phi : X \rightarrow H, \quad x \mapsto K_x \tag{34}$$

that, having the applications in mind, unfortunately is not necessarily injective:

Lemma 2.2 *The map ϕ is injective if and only if for every pair of distinct points $x_1, x_2 \in X$ there exists an $h \in H$ such that $h(x_1) \neq h(x_2)$.*

This follows immediately from the equation $e_x = \langle \cdot, K_x \rangle$.

Utilizing the map ϕ one arrives at a key identity with respect to applications, namely

$$\forall x_1, x_2 \in X : \langle \phi(x_1), \phi(x_2) \rangle = K(x_2, x_1). \quad (35)$$

It follows directly from the definitions of K and ϕ .

Although the family $(K_x)_{x \in X}$ in general does not form a basis of H it generates H in the topological sense:

Proposition 2.3 *For a Hilbert space H functional over X and possessing continuous evaluation functionals the subspace*

$$U := \sum_{x \in X} \mathbb{K}K_x$$

lies dense in H .

Proof. It suffices to prove that only the zero-function is perpendicular to every function K_x . Indeed $\langle h, K_x \rangle = 0$ for all $x \in X$ by (31) is equivalent to $h = 0$. \square

As a consequence of Proposition 2.3 the inclusion $H \subseteq \text{Map}(X, \mathbb{K})$ can be described using the functions $u \in U$ (and thus the functions K_x) only: every function $h \in H$ can be expressed as a limit $\lim_{k \rightarrow \infty} u_k$, $u_k \in U$. The continuity of the evaluation functionals (30) thus yields

$$h(x) = \lim_{k \rightarrow \infty} u_k(x). \quad (36)$$

This fact can be used to reduce analytic to algebraic statements.

2.2 Kernel functions

Let X be a set. Motivated by Proposition 2.1 one calls a function

$$K : X \times X \rightarrow \mathbb{K}$$

with the properties 1 and 2 stated in that proposition a **kernel function (on X)**. Slightly abusing language property 1 is called **symmetry of K** while property 2 is named **positive semi-definiteness**. If for all $n \in \mathbb{N}$ and for pairwise distinct $x_1, \dots, x_n \in X$ the matrices $K(x_1, \dots, x_n)$ are positive definite, then K is called **positive definite**.

The set of kernel functions on X to the field $\mathbb{K} \in \{\mathbb{R}, \mathbb{C}\}$ carries a rich algebraic and order structure that can be utilized effectively in applications. Again the focus in the present section is put onto the facts relevant for the applications in discriminant analysis.

Theorem 2.4 *The set $\mathcal{K}(X)$ of kernel functions on X is closed under pointwise addition, multiplication and scalar multiplication with non-negative scalars $c \in \mathbb{R}$, that is it forms a (linear) cone. In particular $\mathcal{K}(X)$ forms a commutative semi-ring taking pointwise addition and multiplication as compositions.*

Proof. Once the closedness of $\mathcal{K}(X)$ under the operations mentioned in the theorem is verified, the validity of all other ring axioms is obvious since the operations are all defined pointwise.

Addition: for a family $(x_1, \dots, x_n) \in X^n$, $K_1, K_2 \in \mathcal{K}(X)$ and $K := K_1 + K_2$ one has

$$K_1(x_1, \dots, x_n) + K_2(x_1, \dots, x_n) = K(x_1, \dots, x_n).$$

Hence for $z \in \mathbb{K}^n$ one gets

$$z^t K(x_1, \dots, x_n) \bar{z} = z^t K_1(x_1, \dots, x_n) \bar{z} + z^t K_2(x_1, \dots, x_n) \bar{z} \geq 0.$$

Scalar multiplication: for a family $(x_1, \dots, x_n) \in X^n$, $K \in \mathcal{K}(X)$ and positive $c \in \mathbb{R}$ one has

$$z^t (cK(x_1, \dots, x_n)) \bar{z} = c(z^t K(x_1, \dots, x_n) \bar{z}) \geq 0.$$

Multiplication: the proof is based on considering the tensor product of bi-/sesquilinear forms. using the fact that if $\mathcal{B} := (b_1, \dots, b_n)$ is a basis of V , then $(b_i \otimes b_j \mid i, j \in \{1, \dots, n\})$ is a basis of $V \otimes V$, for bi-/sesquilinear forms α, β on some vector space V of finite dimension one can define the tensor product

$$\alpha \otimes \beta : (V \otimes V) \times (V \otimes V) \rightarrow \mathbb{K}$$

through setting

$$(\alpha \otimes \beta)((v_1 \otimes v_2), (w_1 \otimes w_2)) := \alpha(v_1, w_1) \cdot \beta(v_2, w_2). \quad (37)$$

$\alpha \otimes \beta$ is a bi-/sesquilinear form that is symmetric (and positive semi-definite) if α and β are symmetric (and positive semi-definite).

From now on assume that α and β are symmetric and positive semi-definite; then $(\alpha \otimes \beta)|_{U \times U}$ is symmetric and positive semi-definite for every sub-vector space $U \subseteq V \otimes V$. Take U to be the subspace spanned by the symmetric tensors $\mathcal{C} := (b_i \otimes b_i \mid i = 1, \dots, n)$; note that \mathcal{C} is a basis of U .

Let $A = (a_{ij})_{i,j \in \{1, \dots, n\}}$ and $B = (b_{ij})_{i,j \in \{1, \dots, n\}}$ be the matrices of α and β with respect to the basis \mathcal{B} . It is then straightforward to check that the matrix $C \in \mathbb{K}^{n \times n}$ of $(\alpha \otimes \beta)|_{U \times U}$ with respect to the basis \mathcal{C} of U looks like

$$C = \begin{pmatrix} a_{11}b_{11} & a_{12}b_{12} & \dots & a_{1n}b_{1n} \\ a_{21}b_{21} & a_{22}b_{22} & \dots & a_{2n}b_{2n} \\ \cdot & \cdot & \dots & \cdot \\ \cdot & \cdot & \dots & \cdot \\ a_{n1}b_{n1} & a_{n2}b_{n2} & \dots & a_{nn}b_{nn} \end{pmatrix}. \quad (38)$$

Consequently the matrix C is symmetric and positive semi-definite. To denote the particular relationship between A , B and C one writes $C = A \odot B$. The matrix C is frequently called the **Hadamard or Schur product** of A and B . One has thus shown that the Hadamard product of symmetric, positive semi-definite matrices is symmetric and positive semi-definite.

To prove closedness under pointwise multiplication take $(x_1, \dots, x_n) \in X^n$, $K_1, K_2 \in \mathcal{K}(X)$ and let $K := K_1 \cdot K_2$. By definition one then has

$$K(x_1, \dots, x_n) = K_1(x_1, \dots, x_n) \odot K_2(x_1, \dots, x_n),$$

hence positive semi-definiteness of $K(x_1, \dots, x_n)$. The symmetry of K is obvious. □

From the previous proof one can extract the following

Corollary 2.5 *The set $\text{SPSD}(n)$ of symmetric, positive semi-definite $n \times n$ -matrices becomes a commutative semi-ring with 1 if one takes the ordinary matrix addition as addition and the Hadamard product as multiplication.*

For a finite set $X = \{x_1, \dots, x_n\}$ with n elements the map

$$\mathcal{K}(X) \rightarrow \text{SPSD}(n), K \mapsto K(x_1, \dots, x_n)$$

is an isomorphism of semi-rings.

In the context of Corollary 2.5 note that if $K(x_1, \dots, x_n)$ is positive semi-definite, then for every permutation $(x_{s(1)}, \dots, x_{s(n)})$ of the elements x_k and for every subfamily $(x_{k_1}, \dots, x_{k_r})$ the matrices $K(x_{s(1)}, \dots, x_{s(n)})$, $K(x_{k_1}, \dots, x_{k_r})$ are positive semi-definite.

Corollary 2.6 *Let K be a kernel function on X and let $p = \sum_{k=0}^d a_k X^k \in \mathbb{R}[X]$ be a polynomial with positive coefficients, then the function*

$$p(K) = \sum_{k=0}^d a_k K^k$$

understood pointwise is a kernel function on X .

Note that $K^0 := 1$, the function $X \times X \rightarrow \mathbb{K}$ that maps everything to 1. Taking the standard scalar product $\langle \cdot, \cdot \rangle$ for K in Corollary 2.6 yields the widely used **polynomial kernel of degree d** :

$$K(x_1, x_2) := (\langle x_1, x_2 \rangle + c)^d, \quad c > 0, \quad d \in \mathbb{N}. \quad (39)$$

It is not difficult to replace the polynomial in Corollary 2.6 by a power series:

Proposition 2.7 Let $(K_i)_{i \in \mathbb{N}}$ be a sequence of kernel functions on X such that the limit

$$\lim_{i \rightarrow \infty} K_i(x_1, x_2)$$

exists for all $x_1, x_2 \in X$. Then the pointwise limit $K := \lim_{i \rightarrow \infty} K_i$ is a kernel function on X .

Proof. Symmetry of K is straightforward to check. For $(x_1, \dots, x_n) \in X^n$ and $z \in \mathbb{K}^n$ one gets

$$z^t K(x_1, \dots, x_n) \bar{z} = \lim_{i \rightarrow \infty} z^t K_i(x_1, \dots, x_n) \bar{z} \geq 0,$$

since the entries in the matrix $K(x_1, \dots, x_n)$ are the limits of the corresponding entries in the matrices $K_i(x_1, \dots, x_n)$. \square

Corollary 2.8 Let $f : U \rightarrow \mathbb{R}$, $U \subseteq \mathbb{K}$, be a function defined through a (convergent) power series with positive coefficients:

$$f(u) = \sum_{i=0}^{\infty} a_i u^i.$$

Let K be a kernel function on X such that $K(x_1, x_2) \in U$ for all $x_1, x_2 \in X$. Then the pointwise limit $f(K)$ defined through

$$f(K)(x_1, x_2) := \sum_{i=0}^{\infty} a_i K(x_1, x_2)^i$$

is a kernel function on X .

Proof. By Proposition 2.6 the functions

$$K_j := \sum_{i=0}^j a_i K^i$$

are kernel functions. By assumption $f(K) = \lim_{j \rightarrow \infty} K_j$, hence Proposition 2.7 yields the assertion. \square

Corollary 2.8 yields the fact that Gauß' error function is a kernel function. The subsequent result is used in the proof but is also relevant in other contexts:

Proposition 2.9 For every $K \in \mathcal{K}(X)$ and every non-zero function $f : X \rightarrow \mathbb{K}$ the map

$$K' : X \times X \rightarrow \mathbb{K}, (x_1, x_2) \mapsto f(x_1) K(x_1, x_2) \overline{f(x_2)}$$

is a kernel function on X .

Proof. Symmetry is immediate to check. For $(x_1, \dots, x_n) \subseteq X^n$ and $z \in \mathbb{K}^m$ one gets

$$\begin{aligned} z^t K'(x_1, \dots, x_n) \bar{z} &= \sum_{i=1}^n \sum_{j=1}^n z_i f(x_i) K(x_i, x_j) \overline{f(x_j) z_j} \\ &= w^t K(x_1, \dots, x_n) \bar{w} \geq 0, \end{aligned}$$

where $w = (z_1 f(x_1), \dots, z_n f(x_n))$. □

Corollary 2.10 For every $h \in \mathbb{R}$ the function

$$K(x_1, x_2) := e^{-\frac{\|x_1 - x_2\|^2}{h^2}}, \quad (40)$$

is a kernel function on \mathbb{K}^m . It is called the **Gauß kernel of bandwidth h** .

Proof. One first has to rewrite K in an appropriate way – the naive approach of applying Corollary 2.8 directly does not work since the exponent $-\frac{\|x_1 - x_2\|^2}{h^2}$ is not kernel function:

$$e^{-\frac{\|x_1 - x_2\|^2}{h^2}} = [e^{-\frac{\langle x_1, x_1 \rangle}{h^2}} e^{-\frac{\langle x_2, x_2 \rangle}{h^2}}] (e^{\frac{\langle x_1, x_2 \rangle}{h^2}})^2. \quad (41)$$

The factor in squared brackets on the right side of (41) is a kernel function according to Proposition 2.9. Since the scalar product of \mathbb{K}^m is a kernel function Corollary 2.8 yields that $e^{\frac{\langle x_1, x_2 \rangle}{h^2}}$ is a kernel function. Consequently K itself is a kernel function due to Theorem 2.4. □

2.3 The Theorem of Aronszajn-Moore

A functional Hilbert space H having continuous evaluation functionals gives rise to a kernel function K , namely the reproducing kernel of H . From a theoretical point of view it is natural to ask whether every kernel function K arises as the reproducing kernel of some Hilbert space $H(K)$. At the same time the positive answer to this question forms the basis for many applications of kernel functions:

Theorem 2.11 (N. Aronszajn, R. L. Moore) For every kernel function $K : X \times X \rightarrow \mathbb{K}$ there exists a Hilbert space H functional over X and possessing continuous evaluation functionals, such that K is the reproducing kernel of H . This Hilbert space $H = H(K)$ is uniquely determined by K .

Proof. Let $K_y := K(\cdot, y) \in \text{Map}(X, \mathbb{K})$ and consider the subvector space $U \subseteq \text{Map}(X, \mathbb{K})$ generated by the family $(K_y \mid y \in X)$. On U one can define a

symmetric bi- respectively sesquilinear form through

$$\beta : U \times U \rightarrow \mathbb{K}, \left(\sum_{x \in X} a_x K_x, \sum_{y \in X} b_y K_y \right) \mapsto \sum_{x \in X} \sum_{y \in X} a_x K(y, x) \overline{b_y}. \quad (42)$$

Since the family $(K_y \mid y \in X)$ in general need not be linearly independent it remains to show that β is well-defined: to this end one has to verify that for every linear combination

$$u := \sum_{x \in X} a_x K_x = 0$$

and every function $K_y, y \in X$, the equation

$$\sum_{x \in X} a_x K(y, x) = 0$$

holds and similarly for switched roles of the two variables of β . Indeed

$$\sum_{x \in X} a_x K(y, x) = u(y) = 0$$

by assumption about u .

Claim: β is positive definite.

β is positive semi-definite since K is a kernel function. Hence the assertion follows from the Cauchy-Schwarz inequality $|\beta(u_1, u_2)|^2 \leq \beta(u_1, u_1)\beta(u_2, u_2)$ once one knows that $\beta(u_1, u_2) = 0$ for all $u_2 \in U$ implies $u_1 = 0$. Indeed $0 = \beta(u_1, K_y)$ for all $y \in X$ implies $u_1(y) = 0$ for all $y \in X$.

Let H be the completion of U with respect to the norm $\|u\|^2 := \beta(u, u)$. It is well-known that H is a Hilbert space taking the unique continuous extension of β to H as the scalar product.

Claim: H is a subvector space of $\text{Map}(X, \mathbb{K})$.

Express $h \in H$ as a limit $h = \lim_{i \rightarrow \infty} u_i, u_i \in U$. For every $x \in X$ the Cauchy-Schwarz inequality then yields

$$|u_i(x) - u_j(x)|^2 = |\beta(u_i - u_j, K_x)|^2 \leq \beta(u_i - u_j, u_i - u_j)K(x, x), \quad (43)$$

hence $(u_i(x))_{i \in \mathbb{N}}$ is a Cauchy sequence in \mathbb{K} . If $h = \lim_{i \rightarrow \infty} v_i, v_i \in U$, is a second representation of h , then $(u_i - v_i)_{i \in \mathbb{N}}$ is a zero-sequence. Consequently by (43) applied to $u_i - v_i$ the sequence $(u_i(x) - v_i(x))_{i \in \mathbb{N}}$ is a zero-sequence too. Therefore one can interpret h as a function $X \rightarrow \mathbb{K}$ through setting

$$h(x) := \lim_{i \rightarrow \infty} u_i(x).$$

This interpretation leads to a linear map $H \rightarrow \text{Map}(X, \mathbb{K})$.

Claim: K is the reproducing kernel of H .

Indeed by definition of β for every $x \in X$ one has

$$\begin{aligned} e_x(h) = h(x) &= \lim_{i \rightarrow \infty} u_i(x) \\ &= \lim_{i \rightarrow \infty} \beta(u_i, K_x) \\ &= \beta(\lim_{i \rightarrow \infty} u_i, K_x) \\ &= \beta(h, K_x). \end{aligned}$$

Note that this also shows that e_x is continuous.

Finally assume that H_i , $i = 1, 2$, are functional Hilbert spaces both possessing $K : X \times X \rightarrow \mathbb{K}$ as their reproducing kernel. Let U_i be the subvector space of H_i spanned by the functions $K_x = K(\cdot, x)$. Note that for $h \in U_i$ one has

$$h(x) = \sum_{j=1}^r a_{x_j} K(x, x_j) \text{ so that } U_1 = U_2 \text{ as subvector spaces of } \text{Map}(X, \mathbb{K})$$

holds. Similarly for the restrictions $(\|\cdot\|_i)|_{U_i}$ of the norms of the H_i one deduces

$$\|h\|_1^2 = \sum_{j=1}^r \sum_{k=1}^r a_{x_j} K(x_k, x_j) \overline{a_{x_k}} = \|h\|_2^2.$$

Consequently using the remark (36) following Proposition 2.3 one concludes $H_1 = H_2$. \square

2.4 Dimension

For a thorough analysis of the results one gets by using Kernel Fisher discriminant functions it is necessary to have some idea about the dimension of the Hilbert space $H(K)$ associated to a kernel function $K : X \times X \rightarrow \mathbb{K}$. Since $H(K)$ is functional over X one gets

$$\dim H(K) \leq |X| \tag{44}$$

if X is a finite set. Moreover as a related general result one should mention:

Proposition 2.12 *For a kernel function $K \in \mathcal{K}(X)$ the functions $(K_x \mid x \in X)$ are \mathbb{K} -linearly independent if and only if K is positive definite.*

Proof. The linear relation

$$\sum_{i=1}^n a_i K_{x_i} = 0$$

is equivalent with

$$\begin{aligned} 0 &= \left\langle \sum_{i=1}^n a_i K_{x_i}, \sum_{j=1}^n a_j K_{x_j} \right\rangle \\ &= \sum_{i=1}^n \sum_{j=1}^n a_i \overline{a_j} K(x_j, x_i). \end{aligned}$$

□

Corollary 2.13 *For a positive definite kernel function $K \in \mathcal{K}(X)$ the Hilbert space $H(K)$ has finite dimension if and only if X is finite; then the family $(K_x \mid x \in X)$ forms a basis of $H(K)$.*

The proof of Proposition 2.12 yields a criterion for the injectivity of the map $\phi : X \rightarrow H(K)$ – a fundamental property in the context of discriminant functions.

Corollary 2.14 *If the kernel function $K \in \mathcal{K}(X)$ has the property*

$$\forall x_1, x_2 \in X, x_1 \neq x_2 : K(x_1, x_2) \text{ is positive definite,}$$

then $H(K)$ separates the points of X and thus ϕ is injective.

Proof. By the argument used in the proof of Proposition 2.12 the functions K_{x_1}, K_{x_2} for $x_1 \neq x_2$ are linearly independent hence $K_{x_1} \neq K_{x_2}$. □

For a polynomial kernel the functions $(K_x \mid x \in X)$ can be linearly dependent as the case of degree one shows. Moreover the dimension of $H(K)$ depends on X and on the degree of the kernel. For »large« X one gets:

Proposition 2.15 *For a polynomial kernel $K(x_1, x_2) := (\langle x_1, x_2 \rangle + c)^d$ the following properties are valid:*

- (1) *if X contains a non-empty open set of \mathbb{K}^m , then the map $\phi : X \rightarrow H(K), x \mapsto K_x$ is injective,*
- (2) $\dim H(K) \leq \binom{m+d}{d}$ *with equality for $X = \mathbb{K}^m$.*

Proof. 1.: Assume that $K_{x_1} = K_{x_2}$ on X for some $x_1, x_2 \in X$. Then

$$\left(\frac{\langle x, x_1 \rangle + c}{\langle x, x_2 \rangle + c} \right)^d = 1,$$

for all $x \in X$ at which the rational function on the left-hand side is defined; denote that set by U . Since X contains a non-empty open set and $\{x \in \mathbb{K}^m \mid \langle x, x_2 \rangle + c = 0\}$ is a hyperplane in \mathbb{K}^m , the set U contains a

non-empty open set of \mathbb{K}^m . On U the rational function

$$f(x) := \frac{\langle x, x_1 \rangle + c}{\langle x, x_2 \rangle + c}$$

attains only finitely many values, namely a subset of the d -th roots of unity. Consequently f is constant, that is $\langle x, x_1 \rangle + c = a(\langle x, x_2 \rangle + c)$, which implies $a = 1$ and thus $x_1 = x_2$ as asserted.

2.: The Hilbert space $H(K)$ is a subvector space of the vector space $P(X, d)$ of polynomial functions on X in m variables and of degree d . The restriction yields a surjective linear map $P(d) \rightarrow P(X, d)$, where $P(d)$ denotes the vector space of polynomial functions on \mathbb{K}^m in m variables and of degree d . Therefore the well-known formula $\dim P(d) = \binom{m+d}{d}$ completes the proof. \square

Gauß kernels behave much nicer than polynomial kernels:

Lemma 2.16 For pairwise distinct numbers $x_1, \dots, x_n \in \mathbb{C}$ the functions

$$e^{-\|x-x_k\|^2}, \quad k = 1, \dots, n,$$

are linearly independent over \mathbb{C} .

Proof. Take a linear relation

$$\sum_{k=1}^n a_k e^{-\|x-x_k\|^2} = 0, \quad a_k \in \mathbb{C},$$

between functions as stated in the lemma and rewrite it in the form

$$0 = \sum_{k=1}^n (a_k e^{-\|x_k\|^2}) e^{-\|x\|^2} e^{2\operatorname{Re}\langle x, x_k \rangle} = \sum_{k=1}^n b_k e^{-\|x\|^2} e^{2\operatorname{Re}\langle x, x_k \rangle},$$

where $b_k := a_k e^{-\|x_k\|^2}$; then:

$$\sum_{k=1}^n b_k e^{2\operatorname{Re}\langle x, x_k \rangle} = 0$$

and since this equation holds for all $x \in \mathbb{R}^m$ one gets the homogenous system

$$\sum_{k=1}^n b_k e^{2\operatorname{Re}\langle jx, x_k \rangle} = 0, \quad j = 0, \dots, n-1, \quad (45)$$

for fixed $x \in \mathbb{R}^m$. Next choose $x \in \mathbb{R}^m$ such that the values $\operatorname{Re} \langle x, x_k \rangle$, $k = 1, \dots, n$, are pairwise distinct. Then the coefficient matrix

$$A := \begin{pmatrix} 1 & 1 & \dots & 1 \\ e^{2\operatorname{Re} \langle x, x_1 \rangle} & e^{2\operatorname{Re} \langle x, x_2 \rangle} & \dots & e^{2\operatorname{Re} \langle x, x_n \rangle} \\ e^{2\operatorname{Re} \langle 2x, x_1 \rangle} & e^{2\operatorname{Re} \langle 2x, x_2 \rangle} & \dots & e^{2\operatorname{Re} \langle 2x, x_n \rangle} \\ \vdots & \vdots & \dots & \vdots \\ e^{2\operatorname{Re} \langle (n-1)x, x_1 \rangle} & e^{2\operatorname{Re} \langle (n-1)x, x_2 \rangle} & \dots & e^{2\operatorname{Re} \langle (n-1)x, x_n \rangle} \end{pmatrix}$$

of (45) is a Vandermonde matrix. By the choice of the values $\operatorname{Re} \langle x, x_k \rangle$ and the injectivity of the real exponential function A is invertible. Consequently the system (45) only has the trivial solution $b_1 = \dots = b_n = 0$, which implies $a_1 = \dots = a_n = 0$ thus completing the proof. \square

Lemma 2.16 combined with Proposition 2.12 yields:

Proposition 2.17 *The Gauß kernel $K : X \times X \rightarrow \mathbb{K}$ on some set $X \subseteq \mathbb{K}^m$ is positive definite. Consequently the Hilbert space $H(K)$ has finite dimension if and only if X is finite; then the equation*

$$\dim H(K) = |X|$$

holds.

3 Kernel Fisher discriminant functions

Building on the theoretical basis layed in Section 2 in the present section explicit formulae for kernel Fisher discriminant functions are derived. Next an algorithm to estimate the parameters in these formulae using classified data is described. Finally an example is presented.

3.1 Explicit formulae

Let $X = X_{-1} \cup X_{+1} \subset \mathbb{R}^m$ be a finite set (of attributes) divided into two disjoint groups. Fix a kernel function

$$K : \mathbb{R}^m \times \mathbb{R}^m \rightarrow \mathbb{R}$$

and consider the functional Hilbert space $H(K|_{X \times X})$ along with the associated map

$$X \xrightarrow{\phi} H(K|_{X \times X})$$

defined in (34). Throughout the whole section it is assumed that ϕ is injective; consequently $\phi(X_{-1}) \cap \phi(X_{+1}) = \emptyset$.

The Hilbert space $H(K|_{X \times X})$ has finite dimension (44). Consequently Fisher's approach as formulated in Subsection 1.2 can be used to separate the sets $\phi(X_{-1})$ and $\phi(X_{+1})$ using a hyperplane: note first that due to Proposition 2.3 the family $(\phi(x) \mid x \in X)$ generates the space $H(K|_{X \times X})$. Choosing a numbering $X = \{x_1, \dots, x_n\}$ a Fisher discriminant function

$$d : H(K|_{X \times X}) \rightarrow \mathbb{R}, h \mapsto \langle a^*, h \rangle + c^*, \quad (46)$$

can thus be defined by taking a solution $\alpha^* = (a_1^*, \dots, a_n^*) \in \mathbb{R}^n$ of the optimization problem (25) replacing the elements x_i by the elements $\phi(x_i)$. In that way one arrives at

$$a^* = \sum_{j=1}^n a_j^* \phi(x_j)$$

and

$$c^* = -\frac{1}{2}(\alpha^*)^t(m_{-1} + m_{+1}),$$

where

$$m_k = \frac{1}{|X_k|} \sum_{x \in X_k} (\langle \phi(x_1), \phi(x) \rangle, \dots, \langle \phi(x_n), \phi(x) \rangle) \in \mathbb{R}^n$$

for $k \in \{-1, +1\}$ – see (19) and (21). The matrices B and W in the current setting of the optimization problem (25) are defined through

$$B = (m_{-1} - m_{+1})(m_{-1} - m_{+1})^t$$

and

$$W := KK^t - |X_{-1}|m_{-1}m_{-1}^t - |X_{+1}|m_{+1}m_{+1}^t$$

where

$$K := (\langle \phi x_i, \phi x_j \rangle)_{i,j \in \{1, \dots, n\}} \in \mathbb{R}^{n \times n}, \quad (47)$$

see (23), (26) and (24). At that point the essence of the kernel method becomes clearly visible: the solutions α^* of the optimization problem (25) for the grouping $\phi(X) = \phi(X_{-1}) \cup \phi(X_{+1})$ are determined by quantities – namely the matrices B and W – that themselves depend on scalar products of the form $\langle \phi(x), \phi(x') \rangle$, $x, x' \in X$ only. Hence due to the key identity $\langle \phi(x), \phi(x') \rangle = K(x', x)$ (35) these solutions can be computed working in the original sample space \mathbb{R}^m instead of $H(K|_{X \times X})$. Note that the matrix K defined in (47) is indeed »equal« (in an obvious sense) to the kernel function $K : X \times X \rightarrow \mathbb{R}$.

The discriminant function to separate the groups X_{-1} and X_{+1} is given by $d \circ \phi : X \rightarrow \mathbb{R}$, which can be expressed explicitly using α^* , the kernel function K and the data X : for $x \in X$ we get

$$\begin{aligned} (d \circ \phi)(x) &= \left\langle \sum_{j=1}^n a_j^* \phi(x_j), \phi(x) \right\rangle + c^* \\ &= \sum_{j=1}^n a_j^* K(x, x_j) + c^*. \end{aligned} \quad (48)$$

New samples $x \in \mathbb{R}^m$ are classified by plugging in x into (48) and using the decision function $\text{sign}(d(\phi(x)))$. Formally this is possible because the kernel function K is defined on the whole of \mathbb{R}^m . Conceptually this procedure can be justified in the usual way: if the data set X sufficiently faithfully represents the distributions underlying the groups Ω_{-1} and Ω_{+1} , then a discriminant function derived from these data is supposed to generalize well.

3.2 Computation

The computation of the kernel Fisher discriminant function (48) from the given data and a kernel function requires to handle the in general huge kernel matrix K (47) as well as the derived matrices B and W . One therefore has to carefully choose among the existing methods for solving the optimization problem (25) in the present context. In this subsection an algorithm for the approximative solution of (25) proposed in [Mik3] is described without going into the details or giving proofs.

Reformulation of the optimization problem: given a grouping $X = X_{-1} \cup X_{+1}$ and a kernel function $K \in \mathcal{K}(\mathbb{R}^m)$ in the article [Mik2] the problem of finding a discriminant function on \mathbb{R}^m of the form

$$d(x) = \sum_{j=1}^n a_j K(x, x_j) + c \quad (49)$$

is considered as a regression problem for the label function $\ell : X \rightarrow \{-1, +1\}$. Let $X = \{x_1, \dots, x_n\}$ be some numbering of X and define

$$\xi(\alpha, c) := (d(x_j) - \ell(x_j))_{j=1, \dots, n}$$

to be the vector of deviations of the discriminant values $d(x_j)$ from the labels; here $\alpha = (a_1, \dots, a_n)$. Moreover let

$$y := (\ell(x_j))_{j=1, \dots, n}$$

be the vector of labels, let K be the kernel matrix (47) and define

$$\mathbf{1} := (1, \dots, 1) \in \mathbb{R}^n, \quad \mathbf{1}_{+1} = \frac{1}{2}(y + \mathbf{1}), \quad \mathbf{1}_{-1} = \mathbf{1} - \mathbf{1}_{+1}.$$

Finally let $P : \mathbb{R}^n \rightarrow \mathbb{R}$ be a function that penalizes solutions α with »many« non-zero coordinates with a penalty weight $R \geq 0$. Then according to [Mik2]:

Proposition 3.1 *For every penalty function P and constant $R > 0$ the optimization problems*

$$\min(\|\xi(\alpha, c)\|^2 + RP(\alpha) \mid \alpha \in \mathbb{R}^n, c \in \mathbb{R}) \quad (50)$$

$$K\alpha + c\mathbf{1} = y + \xi \quad (51)$$

$$\langle \mathbf{1}_i, \xi \rangle = 0, \quad i \in \{-1, +1\} \quad (52)$$

and

$$\min(\alpha^t W \alpha + RP(\alpha) \mid \alpha \in \mathbb{R}^n) \quad (53)$$

$$\langle \alpha, (m_{+1} - m_{-1}) \rangle = 2 \quad (54)$$

have the same solutions. For the second optimization problem the constant c is computed using the formula (28).

Note that the second optimization problem is just a reformulation of (25): the fraction

$$\frac{\alpha^t B \alpha}{\alpha^t W \alpha}$$

does not change its value when replacing α by a multiple $\lambda \alpha$, $\lambda \neq 0$. This substitution corresponds to multiplying the associated discriminant function by the factor λ . We can thus assume (54) and consequently that $\alpha^t B \alpha$ has a fixed

(positive) value, which in turn yields that it suffices to minimize the denominator $\alpha^t W \alpha$. In that way one arrives at the second optimization problem appearing in Proposition 3.1. Taking into account that the matrix W can and frequently will be singular, the penalty term $RP(\alpha)$ can also be understood as a form of regularization.

A greedy algorithm ... to solve the optimization problem (50) and thus (53) approximatively. It is proposed in the article [Mik3] and is based on the idea of building up the discriminant function (48) stepwise by adding one suitable term $K(x, x_i)$ at each step and stopping the process once the performance of the discriminant function is sufficient. In this way a sparse approximate solution of (50) can be efficiently computed. In principal, choosing the number of non-zero coordinates of α as the penalty function $P(\alpha)$, sparse solutions could also be obtained avoiding the stepwise approach but their computation is not efficient. In the greedy algorithm presented here the penalty term $RP(\alpha)$ should better be considered as a regularization.

To give an explicit description of the crucial step of adding a new term to the discriminant function assume that an approximative solution of (50) involving only $r \geq 1$ terms has already been determined and let $I = \{i_1, \dots, i_r\}$ be the indices of the samples $x_i \in X$ such that $K(x, x_i)$ appears in this solution. Indeed only the terms $K(x, x_i)$, $i \in I$, are of interest here. The associated coefficients a_i will be recomputed at each step.

Take $i_{r+1} \in \{1, \dots, n\} \setminus I$ and set $I' := I \cup \{i_{r+1}\}$. The aim now is to efficiently compute an approximative solution of (50) involving the $r + 1$ terms $K(x, x_i)$, $i \in I'$. To this end one rewrites (50), (51) and (52) in matrix form. Setting

$$\beta := (c, a_{i_1}, \dots, a_{i_{r+1}}) \in \mathbb{R}^{r+2}$$

and

$$K_{r+1} := \begin{pmatrix} K(x_1, x_{i_1}) & K(x_1, x_{i_2}) & \dots & K(x_1, x_{i_{r+1}}) \\ K(x_2, x_{i_1}) & K(x_2, x_{i_2}) & \dots & K(x_2, x_{i_{r+1}}) \\ \cdot & \cdot & \dots & \cdot \\ \cdot & \cdot & \dots & \cdot \\ K(x_n, x_{i_1}) & K(x_n, x_{i_2}) & \dots & K(x_n, x_{i_{r+1}}) \end{pmatrix} \in \mathbb{R}^{n \times (r+1)}$$

the matrix form of the considered optimization problem is

$$\min(\beta^t H \beta - \frac{1}{2} \gamma^t \beta + n \quad | \quad \beta \in \mathbb{R}^{r+2}) \quad (55)$$

$$A_{-1}^t \beta + |X_{-1}| = 0 \quad (56)$$

$$A_{+1}^t \beta - |X_{+1}| = 0, \quad (57)$$

where

$$\begin{aligned}\gamma &:= (|X_{+1}| - |X_{-1}|, K_{r+1}^t y) \in \mathbb{R}^{r+2}, \\ A_k &:= (|X_k|, K_{r+1}^t \mathbf{1}_k) \in \mathbb{R}^{r+2}, \quad k \in \{-1, +1\}, \\ H &:= \begin{pmatrix} n & \mathbf{1}^t K_{r+1} \\ K_{r+1}^t \mathbf{1} & K_{r+1}^t K_{r+1} + R \end{pmatrix} \in \mathbb{R}^{(r+2) \times (r+2)}.\end{aligned}$$

The optimization problem (55), (56) and (57) can be solved using Lagrange multipliers thus obtaining:

$$\beta^* = H^{-1}(\gamma - \lambda_{+1}^* A_{+1} - \lambda_{-1}^* A_{-1}), \quad (58)$$

where the multipliers $(\lambda_{-1}^*, \lambda_{+1}^*)$ form a solution of the optimization problem

$$\max\left(-\frac{1}{2}\lambda^t Q \lambda + L\lambda - \frac{1}{2}\gamma^t H^{-1}\gamma + \frac{n}{2} \mid \lambda = (\lambda_{+1}, \lambda_{-1}) \in \mathbb{R}\right), \quad (59)$$

where

$$Q := \begin{pmatrix} A_{+1}^t H^{-1} A_{+1} & A_{+1}^t H^{-1} A_{-1} \\ A_{-1}^t H^{-1} A_{+1} & A_{-1}^t H^{-1} A_{-1} \end{pmatrix} \in \mathbb{R}^{2 \times 2},$$

and

$$L := ((-|X_{+1}| + \gamma^t H^{-1} A_{+1}) \quad (|X_{-1}| + \gamma^t H^{-1} A_{-1})) \in \mathbb{R}^2.$$

Note that the optimization problem (59) can be solved analytically by taking the first derivative of the quadratic function involved.

Note further that due to the particular form of the matrices H one has an equation of the form

$$H_{r+1} = \begin{pmatrix} H_r & v \\ v^t & s \end{pmatrix}, \quad v \in \mathbb{R}^{r+1}, \quad s \in \mathbb{R},$$

relating the matrix H for r terms in the discriminant function with the matrix for $r + 1$ terms. Consequently the inverse H^{-1} for $r + 1$ terms can be computed efficiently from the inverse for r terms using the well-known Sherman-Woodbury-formula.

The algorithm can now be formulated as follows:

- (1) Start with the empty set $I = \emptyset$ of indices $i \in \{1, \dots, n\}$ such that $K(x, x_i)$ appears in the discriminant function.
- (2) Compute the solutions $\beta^*(i')$ for all index sets $I \cup \{i'\}$, $i' \in \{1, \dots, n\} \setminus I$, using the formula (58).
- (3) Select an index $i_0 \in \{1, \dots, n\} \setminus I$ such that the objective function (55) at the solution $\beta^*(i_0)$ is minimal among all of the objective function values at the solutions $\beta^*(i)$. Set $I := I \cup \{i_0\}$.

- (4) Stop if $|I|$ exceeds a predefined maximal number of terms in the discriminant function or if the change of the value of the objective function (55) compared to the previous addition of a new term lies below a predefined bound. Otherwise go to step 2.

3.3 An example

In the sequel the results of an application of the greedy algorithm described in Subsection 3.2 to an artificial two-dimensional data set are presented with the aim to give a visual impression of Kernel Fisher discriminant functions in a concrete case (and not of the performance of the algorithm used).

The data set $X = X_{-1} \cup X_{+1}$ shown in Figure 1 consists of 1640 samples $x \in \mathbb{R}^2$, each of the groups X_{-1} and X_{+1} having 820 elements. It has been created by randomly distributing points about two semi-circles using a normal distribution with standard deviation $\sigma = 0.8$. The »center points« of the semi-circles lie at $(-0.5, 0)$ and $(0, 0.5)$, both having a »radius« equal to 1.




Figure 1 data set

First a Gauß kernel $K(x_1, x_2) = e^{-\frac{\|x_1 - x_2\|^2}{h^2}}$ (40) having bandwidth $h = 0.858$ was used. The value of h has been estimated from the data using 5-fold cross validation – knowing the way the data set has been created a reasonable result.


The algorithm gives the approximate discriminant function

$$d_{\text{Gau\ss}}(x) := \begin{aligned} & -0.7591 \cdot K(x, x_1) - 2.8579 \cdot K(x, x_2) - 2.4329 \cdot K(x, x_3) \\ & - 2.0455 \cdot K(x, x_4) - 1.6439 \cdot K(x, x_5) + 1.3246, \end{aligned} \quad (60)$$

with

$$\begin{aligned} x_1 &= (-0.5256, 1.0841), & x_2 &= (-1.9011, 0.1797), \\ x_3 &= (0.7500, 0.2981), & x_4 &= (-1.1620, 1.6343), \\ x_5 &= (0.0948, 1.5043). \end{aligned}$$

Figure 2 shows various level sets $d_{\text{Gau\ss}}^{-1}(l)$, $l \in \mathbb{R}$, of the function (60); the level curve $d_{\text{Gau\ss}}^{-1}(0)$, that is the separating curve, is drawn in black colour. In addition the level of the function values $d_{\text{Gau\ss}}(x)$ is depicted through a yellow-to-orange colour scale symbolising decreasing values of $d_{\text{Gau\ss}}(x)$. The locations of the points x_1, \dots, x_5 are marked as white points.



level curves of $d_{\text{Gau\ss}}(x)$

Figure 2

Using the classification rule $\ell(x) = \text{sgn}(d_{\text{Gau\ss}}(x))$ (11) yields misclassification rates of 3.9% in the group X_{-1} and 1.9% in the group X_{+1} .

In a second run a polynomial kernel $K(x_1, x_2) = (\langle x_1, x_2 \rangle + c)^3$ (39) of degree 3 with constant term $c = 0.75$ has been used, the value of c again being estimated using 5-fold crossvalidation. The resulting estimated discriminant function is:

$$d_{\text{poly}}(x) := \begin{aligned} & -7.1287 \cdot K(x, x_1) + 0.3231 \cdot K(x, x_2) + 0.3612 \cdot K(x, x_3) \\ & - 0.5870 \cdot K(x, x_4) + 0.9951 \cdot K(x, x_5) + 2.4521, \end{aligned} \quad (61)$$

with

$$\begin{aligned} x_1 &= (-0.0150, 0.1517), & x_2 &= (0.0049, 0.7557), \\ x_3 &= (1.8009, 0.2819), & x_4 &= (1.4297, 0.2090), \\ x_5 &= (-0.2253, 0.0755). \end{aligned}$$

Figure 3 depicts the level sets of d_{poly} , the separating curve and the positions of the x_k using the same symbolism as in the previous example.

The classification rule $\ell(x) = \text{sgn}(d_{\text{poly}}(x))$ this time yields a misclassification rate of 3.9% in the group X_{-1} and 5.0% in the group X_{+1} .





Figure 3 level curves of $d_{\text{poly}}(x)$

Assuming that the functions $d_{\text{Gau}\beta}$ and d_{poly} are good approximations to the respective optimal discriminant function (46), their function values $d_{\text{Gau}\beta}(x)$ and $d_{\text{poly}}(x)$, $x \in X$, up to a constant factor and a translation represent the orthogonal projections of the points $\phi(x)$ to the line $\mathbb{R}\alpha \subseteq H(K|_{X \times X})$, where α is the vector of coefficients of $d_{\text{Gau}\beta}$ respectively d_{poly} amended by zeros.

The left plot in Figure 4 shows a histogram of the distribution of $d_{\text{Gau}\beta}(X_{-1})$ plotted in blue and of $d_{\text{Gau}\beta}(X_{+1})$ plotted in red. Bars in different colour directly beside each other refer to one and the same bin of the histogram. The right plot in Figure 4 shows the distribution of $d_{\text{poly}}(X_{-1})$ and $d_{\text{poly}}(X_{+1})$ using the same colour code.




distribution of $d_{\text{GauB}}(X)$ (left) and $d_{\text{poly}}(X)$ (right)

Figure 4

Figure (5) shows the results of the classical Fisher approach. The discriminant function here is:

$$d_F(x) := \langle (0.4951, -1.4871), x \rangle \quad (62)$$

with a misclassification rate of 10.5% in group X_{-1} and 8.9% in group X_{+1} . Note that one can determine Fisher's linear discriminant function approximately by applying Mika's algorithm utilizing the kernel function $K(x_1, x_2) := \langle x_1, x_2 \rangle$. However here the usual approach has been chosen.



level curves of $d_F(x)$ (left), distribution of $d_F(X)$ (right)

Figure 5

References

- [Aiz] M. Aizerman, E. Braverman, L. Rozonoer *Theoretical foundations of the potential function method in pattern recognition learning*, Automation and Remote Control **25** (1964), 821-837.
- [Aro] N. Aronszajn, *Theory of reproducing kernels*, Trans. Amer. Math. Soc. **68** (1950), 337-404.
- [Fis] R. A. Fisher, *The Use of Multiple Measurements in Taxonomic Problems*, Annals of Eugenics **7** (1936), 179-188.
- [Mik1] S. Mika et. al., *Fisher Discriminant Analysis With Kernels*, Neural Networks for Signal Processing **IX** (1999), 41-48.
- [Mik2] S. Mika et. al., *A Mathematical Programming Approach to the Kernel Fisher Algorithm*, Advances in Neural Information Processing Systems **13** (2001).
- [Mik3] S. Mika et. al., *An Improved Training Algorithm for Kernel Fisher Discriminants*, Proceedings AISTATS 2001.
- [Moo] E. H. Moore, *On properly positive Hermitian matrices*, Bull. Amer. Math. Soc. **23(59)** (1916), 66-67.
- [ST-N] J. Shawe-Taylor, N. Cristianini, *Kernel Methods for Pattern Analysis*, Cambridge University Press, 2004.

Published reports of the Fraunhofer ITWM

The PDF-files of the following reports are available under:

www.itwm.fraunhofer.de/de/zentral__berichte/berichte

1. D. Hietel, K. Steiner, J. Struckmeier
A Finite - Volume Particle Method for Compressible Flows
(19 pages, 1998)
2. M. Feldmann, S. Seibold
Damage Diagnosis of Rotors: Application of Hilbert Transform and Multi-Hypothesis Testing
Keywords: Hilbert transform, damage diagnosis, Kalman filtering, non-linear dynamics
(23 pages, 1998)
3. Y. Ben-Haim, S. Seibold
Robust Reliability of Diagnostic Multi-Hypothesis Algorithms: Application to Rotating Machinery
Keywords: Robust reliability, convex models, Kalman filtering, multi-hypothesis diagnosis, rotating machinery, crack diagnosis
(24 pages, 1998)
4. F.-Th. Lentjes, N. Siedow
Three-dimensional Radiative Heat Transfer in Glass Cooling Processes
(23 pages, 1998)
5. A. Klar, R. Wegener
A hierarchy of models for multilane vehicular traffic
Part I: Modeling
(23 pages, 1998)

Part II: Numerical and stochastic investigations
(17 pages, 1998)
6. A. Klar, N. Siedow
Boundary Layers and Domain Decomposition for Radiative Heat Transfer and Diffusion Equations: Applications to Glass Manufacturing Processes
(24 pages, 1998)
7. I. Choquet
Heterogeneous catalysis modelling and numerical simulation in rarified gas flows
Part I: Coverage locally at equilibrium
(24 pages, 1998)
8. J. Ohser, B. Steinbach, C. Lang
Efficient Texture Analysis of Binary Images
(17 pages, 1998)
9. J. Orlik
Homogenization for viscoelasticity of the integral type with aging and shrinkage
(20 pages, 1998)
10. J. Mohring
Helmholtz Resonators with Large Aperture
(21 pages, 1998)
11. H. W. Hamacher, A. Schöbel
On Center Cycles in Grid Graphs
(15 pages, 1998)
12. H. W. Hamacher, K.-H. Küfer
Inverse radiation therapy planning - a multiple objective optimisation approach
(14 pages, 1999)
13. C. Lang, J. Ohser, R. Hilfer
On the Analysis of Spatial Binary Images
(20 pages, 1999)
14. M. Junk
On the Construction of Discrete Equilibrium Distributions for Kinetic Schemes
(24 pages, 1999)
15. M. Junk, S. V. Raghurame Rao
A new discrete velocity method for Navier-Stokes equations
(20 pages, 1999)
16. H. Neunzert
Mathematics as a Key to Key Technologies
(39 pages (4 PDF-Files), 1999)
17. J. Ohser, K. Sandau
Considerations about the Estimation of the Size Distribution in Wicksell's Corpuscle Problem
(18 pages, 1999)
18. E. Carrizosa, H. W. Hamacher, R. Klein, S. Nickel
Solving nonconvex planar location problems by finite dominating sets
Keywords: Continuous Location, Polyhedral Gauges, Finite Dominating Sets, Approximation, Sandwich Algorithm, Greedy Algorithm
(19 pages, 2000)
19. A. Becker
A Review on Image Distortion Measures
Keywords: Distortion measure, human visual system
(26 pages, 2000)
20. H. W. Hamacher, M. Labbé, S. Nickel, T. Sonneborn
Polyhedral Properties of the Uncapacitated Multiple Allocation Hub Location Problem
Keywords: integer programming, hub location, facility location, valid inequalities, facets, branch and cut
(21 pages, 2000)
21. H. W. Hamacher, A. Schöbel
Design of Zone Tariff Systems in Public Transportation
(30 pages, 2001)
22. D. Hietel, M. Junk, R. Keck, D. Teleaga
The Finite-Volume-Particle Method for Conservation Laws
(16 pages, 2001)
23. T. Bender, H. Hennes, J. Kalcsics, M. T. Melo, S. Nickel
Location Software and Interface with GIS and Supply Chain Management
Keywords: facility location, software development, geographical information systems, supply chain management
(48 pages, 2001)
24. H. W. Hamacher, S. A. Tjandra
Mathematical Modelling of Evacuation Problems: A State of Art
(44 pages, 2001)
25. J. Kuhnert, S. Tiwari
Grid free method for solving the Poisson equation
Keywords: Poisson equation, Least squares method, Grid free method
(19 pages, 2001)
26. T. Götz, H. Rave, D. Reinel-Bitzer, K. Steiner, H. Tiemeier
Simulation of the fiber spinning process
Keywords: Melt spinning, fiber model, Lattice Boltzmann, CFD
(19 pages, 2001)
27. A. Zemitis
On interaction of a liquid film with an obstacle
Keywords: impinging jets, liquid film, models, numerical solution, shape
(22 pages, 2001)
28. I. Ginzburg, K. Steiner
Free surface lattice-Boltzmann method to model the filling of expanding cavities by Bingham Fluids
Keywords: Generalized LBE, free-surface phenomena, interface boundary conditions, filling processes, Bingham viscoplastic model, regularized models
(22 pages, 2001)
29. H. Neunzert
»Denn nichts ist für den Menschen als Menschen etwas wert, was er nicht mit Leidenschaft tun kann«
Vortrag anlässlich der Verleihung des Akademiepreises des Landes Rheinland-Pfalz am 21.11.2001
Keywords: Lehre, Forschung, angewandte Mathematik, Mehrskalalanalyse, Strömungsmechanik
(18 pages, 2001)
30. J. Kuhnert, S. Tiwari
Finite pointset method based on the projection method for simulations of the incompressible Navier-Stokes equations
Keywords: Incompressible Navier-Stokes equations, Meshfree method, Projection method, Particle scheme, Least squares approximation
AMS subject classification: 76D05, 76M28
(25 pages, 2001)
31. R. Korn, M. Krekel
Optimal Portfolios with Fixed Consumption or Income Streams
Keywords: Portfolio optimisation, stochastic control, HJB equation, discretisation of control problems.
(23 pages, 2002)
32. M. Krekel
Optimal portfolios with a loan dependent credit spread
Keywords: Portfolio optimisation, stochastic control, HJB equation, credit spread, log utility, power utility, non-linear wealth dynamics
(25 pages, 2002)
33. J. Ohser, W. Nagel, K. Schladitz
The Euler number of discretized sets – on the choice of adjacency in homogeneous lattices
Keywords: image analysis, Euler number, neighborhood relationships, cuboidal lattice
(32 pages, 2002)

34. I. Ginzburg, K. Steiner
Lattice Boltzmann Model for Free-Surface flow and Its Application to Filling Process in Casting
Keywords: Lattice Boltzmann models; free-surface phenomena; interface boundary conditions; filling processes; injection molding; volume of fluid method; interface boundary conditions; advection-schemes; up-wind-schemes (54 pages, 2002)
35. M. Günther, A. Klar, T. Materne, R. Wegener
Multivalued fundamental diagrams and stop and go waves for continuum traffic equations
Keywords: traffic flow, macroscopic equations, kinetic derivation, multivalued fundamental diagram, stop and go waves, phase transitions (25 pages, 2002)
36. S. Feldmann, P. Lang, D. Prätzel-Wolters
Parameter influence on the zeros of network determinants
Keywords: Networks, Equicofactor matrix polynomials, Realization theory, Matrix perturbation theory (30 pages, 2002)
37. K. Koch, J. Ohser, K. Schladitz
Spectral theory for random closed sets and estimating the covariance via frequency space
Keywords: Random set, Bartlett spectrum, fast Fourier transform, power spectrum (28 pages, 2002)
38. D. d'Humières, I. Ginzburg
Multi-reflection boundary conditions for lattice Boltzmann models
Keywords: lattice Boltzmann equation, boundary conditions, bounce-back rule, Navier-Stokes equation (72 pages, 2002)
39. R. Korn
Elementare Finanzmathematik
Keywords: Finanzmathematik, Aktien, Optionen, Portfolio-Optimierung, Börse, Lehrerweiterbildung, Mathematikunterricht (98 pages, 2002)
40. J. Kallrath, M. C. Müller, S. Nickel
Batch Presorting Problems: Models and Complexity Results
Keywords: Complexity theory, Integer programming, Assignment, Logistics (19 pages, 2002)
41. J. Linn
On the frame-invariant description of the phase space of the Folgar-Tucker equation
Key words: fiber orientation, Folgar-Tucker equation, injection molding (5 pages, 2003)
42. T. Hanne, S. Nickel
A Multi-Objective Evolutionary Algorithm for Scheduling and Inspection Planning in Software Development Projects
Key words: multiple objective programming, project management and scheduling, software development, evolutionary algorithms, efficient set (29 pages, 2003)
43. T. Bortfeld, K.-H. Küfer, M. Monz, A. Scherrer, C. Thieke, H. Trinkaus
Intensity-Modulated Radiotherapy - A Large Scale Multi-Criteria Programming Problem
Keywords: multiple criteria optimization, representative systems of Pareto solutions, adaptive triangulation, clustering and disaggregation techniques, visualization of Pareto solutions, medical physics, external beam radiotherapy planning, intensity modulated radiotherapy (31 pages, 2003)
44. T. Halfmann, T. Wichmann
Overview of Symbolic Methods in Industrial Analog Circuit Design
Keywords: CAD, automated analog circuit design, symbolic analysis, computer algebra, behavioral modeling, system simulation, circuit sizing, macro modeling, differential-algebraic equations, index (17 pages, 2003)
45. S. E. Mikhailov, J. Orlik
Asymptotic Homogenisation in Strength and Fatigue Durability Analysis of Composites
Keywords: multiscale structures, asymptotic homogenization, strength, fatigue, singularity, non-local conditions (14 pages, 2003)
46. P. Domínguez-Marín, P. Hansen, N. Mladenović, S. Nickel
Heuristic Procedures for Solving the Discrete Ordered Median Problem
Keywords: genetic algorithms, variable neighborhood search, discrete facility location (31 pages, 2003)
47. N. Boland, P. Domínguez-Marín, S. Nickel, J. Puerto
Exact Procedures for Solving the Discrete Ordered Median Problem
Keywords: discrete location, Integer programming (41 pages, 2003)
48. S. Feldmann, P. Lang
Padé-like reduction of stable discrete linear systems preserving their stability
Keywords: Discrete linear systems, model reduction, stability, Hankel matrix, Stein equation (16 pages, 2003)
49. J. Kallrath, S. Nickel
A Polynomial Case of the Batch Presorting Problem
Keywords: batch presorting problem, online optimization, competitive analysis, polynomial algorithms, logistics (17 pages, 2003)
50. T. Hanne, H. L. Trinkaus
knowCube for MCDM – Visual and Interactive Support for Multicriteria Decision Making
Key words: Multicriteria decision making, knowledge management, decision support systems, visual interfaces, interactive navigation, real-life applications. (26 pages, 2003)
51. O. Iliev, V. Laptev
On Numerical Simulation of Flow Through Oil Filters
Keywords: oil filters, coupled flow in plain and porous media, Navier-Stokes, Brinkman, numerical simulation (8 pages, 2003)
52. W. Dörfler, O. Iliev, D. Stoyanov, D. Vassileva
On a Multigrid Adaptive Refinement Solver for Saturated Non-Newtonian Flow in Porous Media
Keywords: Nonlinear multigrid, adaptive refinement, Heston model, stochastic volatility, cliquet options (17 pages, 2003)
53. S. Kruse
On the Pricing of Forward Starting Options under Stochastic Volatility
Keywords: Option pricing, forward starting options, Heston model, stochastic volatility, cliquet options (11 pages, 2003)
54. O. Iliev, D. Stoyanov
Multigrid – adaptive local refinement solver for incompressible flows
Keywords: Navier-Stokes equations, incompressible flow, projection-type splitting, SIMPLE, multigrid methods, adaptive local refinement, lid-driven flow in a cavity (37 pages, 2003)
55. V. Starikovicius
The multiphase flow and heat transfer in porous media
Keywords: Two-phase flow in porous media, various formulations, global pressure, multiphase mixture model, numerical simulation (30 pages, 2003)
56. P. Lang, A. Sarishvili, A. Wirsén
Blocked neural networks for knowledge extraction in the software development process
Keywords: Blocked Neural Networks, Nonlinear Regression, Knowledge Extraction, Code Inspection (21 pages, 2003)
57. H. Knaf, P. Lang, S. Zeiser
Diagnosis aiding in Regulation Thermography using Fuzzy Logic
Keywords: fuzzy logic, knowledge representation, expert system (22 pages, 2003)
58. M. T. Melo, S. Nickel, F. Saldanha da Gama
Largescale models for dynamic multi-commodity capacitated facility location
Keywords: supply chain management, strategic planning, dynamic location, modeling (40 pages, 2003)
59. J. Orlik
Homogenization for contact problems with periodically rough surfaces
Keywords: asymptotic homogenization, contact problems (28 pages, 2004)
60. A. Scherrer, K.-H. Küfer, M. Monz, F. Alonso, T. Bortfeld
IMRT planning on adaptive volume structures – a significant advance of computational complexity
Keywords: Intensity-modulated radiation therapy (IMRT), inverse treatment planning, adaptive volume structures, hierarchical clustering, local refinement, adaptive clustering, convex programming, mesh generation, multi-grid methods (24 pages, 2004)

61. D. Kehrwald
Parallel lattice Boltzmann simulation of complex flows
Keywords: Lattice Boltzmann methods, parallel computing, microstructure simulation, virtual material design, pseudo-plastic fluids, liquid composite moulding (12 pages, 2004)
62. O. Iliev, J. Linn, M. Moog, D. Niedziela, V. Starikovicus
On the Performance of Certain Iterative Solvers for Coupled Systems Arising in Discretization of Non-Newtonian Flow Equations
Keywords: Performance of iterative solvers, Preconditioners, Non-Newtonian flow (17 pages, 2004)
63. R. Ciegis, O. Iliev, S. Rief, K. Steiner
On Modelling and Simulation of Different Regimes for Liquid Polymer Moulding
Keywords: Liquid Polymer Moulding, Modelling, Simulation, Infiltration, Front Propagation, non-Newtonian flow in porous media (43 pages, 2004)
64. T. Hanne, H. Neu
Simulating Human Resources in Software Development Processes
Keywords: Human resource modeling, software process, productivity, human factors, learning curve (14 pages, 2004)
65. O. Iliev, A. Mikelic, P. Popov
Fluid structure interaction problems in deformable porous media: Toward permeability of deformable porous media
Keywords: fluid-structure interaction, deformable porous media, upscaling, linear elasticity, stokes, finite elements (28 pages, 2004)
66. F. Gaspar, O. Iliev, F. Lisbona, A. Naumovich, P. Vabishchevich
On numerical solution of 1-D poroelasticity equations in a multilayered domain
Keywords: poroelasticity, multilayered material, finite volume discretization, MAC type grid (41 pages, 2004)
67. J. Ohser, K. Schladitz, K. Koch, M. Nöthe
Diffraction by image processing and its application in materials science
Keywords: porous microstructure, image analysis, random set, fast Fourier transform, power spectrum, Bartlett spectrum (13 pages, 2004)
68. H. Neunzert
Mathematics as a Technology: Challenges for the next 10 Years
Keywords: applied mathematics, technology, modelling, simulation, visualization, optimization, glass processing, spinning processes, fiber-fluid interaction, turbulence effects, topological optimization, multicriteria optimization, Uncertainty and Risk, financial mathematics, Malliavin calculus, Monte-Carlo methods, virtual material design, filtration, bio-informatics, system biology (29 pages, 2004)
69. R. Ewing, O. Iliev, R. Lazarov, A. Naumovich
On convergence of certain finite difference discretizations for 1D poroelasticity interface problems
Keywords: poroelasticity, multilayered material, finite volume discretizations, MAC type grid, error estimates (26 pages, 2004)
70. W. Dörfler, O. Iliev, D. Stoyanov, D. Vassileva
On Efficient Simulation of Non-Newtonian Flow in Saturated Porous Media with a Multigrid Adaptive Refinement Solver
Keywords: Nonlinear multigrid, adaptive refinement, non-Newtonian in porous media (25 pages, 2004)
71. J. Kalcsics, S. Nickel, M. Schröder
Towards a Unified Territory Design Approach – Applications, Algorithms and GIS Integration
Keywords: territory design, political districting, sales territory alignment, optimization algorithms, Geographical Information Systems (40 pages, 2005)
72. K. Schladitz, S. Peters, D. Reinel-Bitzer, A. Wiegmann, J. Ohser
Design of acoustic trim based on geometric modeling and flow simulation for non-woven
Keywords: random system of fibers, Poisson line process, flow resistivity, acoustic absorption, Lattice-Boltzmann method, non-woven (21 pages, 2005)
73. V. Rutka, A. Wiegmann
Explicit Jump Immersed Interface Method for virtual material design of the effective elastic moduli of composite materials
Keywords: virtual material design, explicit jump immersed interface method, effective elastic moduli, composite materials (22 pages, 2005)
74. T. Hanne
Eine Übersicht zum Scheduling von Baustellen
Keywords: Projektplanung, Scheduling, Bauplanung, Bauindustrie (32 pages, 2005)
75. J. Linn
The Folgar-Tucker Model as a Differential Algebraic System for Fiber Orientation Calculation
Keywords: fiber orientation, Folgar-Tucker model, invariants, algebraic constraints, phase space, trace stability (15 pages, 2005)
76. M. Speckert, K. Dreßler, H. Mauch, A. Lion, G. J. Wierda
Simulation eines neuartigen Prüfsystems für Achserproben durch MKS-Modellierung einschließlich Regelung
Keywords: virtual test rig, suspension testing, multi-body simulation, modeling hexapod test rig, optimization of test rig configuration (20 pages, 2005)
77. K.-H. Küfer, M. Monz, A. Scherrer, P. Süß, F. Alonso, A. S. A. Sultan, Th. Bortfeld, D. Craft, Chr. Thieke
Multicriteria optimization in intensity modulated radiotherapy planning
Keywords: multicriteria optimization, extreme solutions, real-time decision making, adaptive approximation schemes, clustering methods, IMRT planning, reverse engineering (51 pages, 2005)
78. S. Amstutz, H. Andrä
A new algorithm for topology optimization using a level-set method
Keywords: shape optimization, topology optimization, topological sensitivity, level-set (22 pages, 2005)
79. N. Ettrich
Generation of surface elevation models for urban drainage simulation
Keywords: Flooding, simulation, urban elevation models, laser scanning (22 pages, 2005)
80. H. Andrä, J. Linn, I. Matei, I. Shklyar, K. Steiner, E. Teichmann
OPTCAST – Entwicklung adäquater Strukturoptimierungsverfahren für Gießereien Technischer Bericht (KURZFASSUNG)
Keywords: Topologieoptimierung, Level-Set-Methode, Gießprozesssimulation, Gießtechnische Restriktionen, CAE-Kette zur Strukturoptimierung (77 pages, 2005)
81. N. Marheineke, R. Wegener
Fiber Dynamics in Turbulent Flows Part I: General Modeling Framework
Keywords: fiber-fluid interaction; Cosserat rod; turbulence modeling; Kolmogorov's energy spectrum; double-velocity correlations; differentiable Gaussian fields (20 pages, 2005)
Part II: Specific Taylor Drag
Keywords: flexible fibers; $k-\epsilon$ turbulence model; fiber-turbulence interaction scales; air drag; random Gaussian aerodynamic force; white noise; stochastic differential equations; ARMA process (18 pages, 2005)
82. C. H. Lampert, O. Wirjadi
An Optimal Non-Orthogonal Separation of the Anisotropic Gaussian Convolution Filter
Keywords: Anisotropic Gaussian filter, linear filtering, orientation space, nD image processing, separable filters (25 pages, 2005)
83. H. Andrä, D. Stoyanov
Error indicators in the parallel finite element solver for linear elasticity DDFEM
Keywords: linear elasticity, finite element method, hierarchical shape functions, domain decomposition, parallel implementation, a posteriori error estimates (21 pages, 2006)
84. M. Schröder, I. Solchenbach
Optimization of Transfer Quality in Regional Public Transit
Keywords: public transit, transfer quality, quadratic assignment problem (16 pages, 2006)
85. A. Naumovich, F. J. Gaspar
On a multigrid solver for the three-dimensional Biot poroelasticity system in multilayered domains
Keywords: poroelasticity, interface problem, multigrid, operator-dependent prolongation (11 pages, 2006)
86. S. Panda, R. Wegener, N. Marheineke
Slender Body Theory for the Dynamics of Curved Viscous Fibers
Keywords: curved viscous fibers; fluid dynamics; Navier-Stokes equations; free boundary value problem; asymptotic expansions; slender body theory (14 pages, 2006)
87. E. Ivanov, H. Andrä, A. Kudryavtsev
Domain Decomposition Approach for Automatic Parallel Generation of Tetrahedral Grids
Key words: Grid Generation, Unstructured Grid, Delaunay Triangulation, Parallel Programming, Domain Decomposition, Load Balancing (18 pages, 2006)

88. S. Tiwari, S. Antonov, D. Hietel, J. Kuhnert, R. Wegener
A Meshfree Method for Simulations of Interactions between Fluids and Flexible Structures
Key words: Meshfree Method, FPM, Fluid Structure Interaction, Sheet of Paper, Dynamical Coupling (16 pages, 2006)
89. R. Ciegis, O. Iliev, V. Starikovicius, K. Steiner
Numerical Algorithms for Solving Problems of Multiphase Flows in Porous Media
Keywords: nonlinear algorithms, finite-volume method, software tools, porous media, flows (16 pages, 2006)
90. D. Niedziela, O. Iliev, A. Latz
On 3D Numerical Simulations of Viscoelastic Fluids
Keywords: non-Newtonian fluids, anisotropic viscosity, integral constitutive equation (18 pages, 2006)
91. A. Winterfeld
Application of general semi-infinite Programming to Lapidary Cutting Problems
Keywords: large scale optimization, nonlinear programming, general semi-infinite optimization, design centering, clustering (26 pages, 2006)
92. J. Orlik, A. Ostrovska
Space-Time Finite Element Approximation and Numerical Solution of Hereditary Linear Viscoelasticity Problems
Keywords: hereditary viscoelasticity; kern approximation by interpolation; space-time finite element approximation, stability and a priori estimate (24 pages, 2006)
93. V. Rutka, A. Wiegmann, H. Andrä
EJIM for Calculation of effective Elastic Moduli in 3D Linear Elasticity
Keywords: Elliptic PDE, linear elasticity, irregular domain, finite differences, fast solvers, effective elastic moduli (24 pages, 2006)
94. A. Wiegmann, A. Zemitis
EJ-HEAT: A Fast Explicit Jump Harmonic Averaging Solver for the Effective Heat Conductivity of Composite Materials
Keywords: Stationary heat equation, effective thermal conductivity, explicit jump, discontinuous coefficients, virtual material design, microstructure simulation, EJ-HEAT (21 pages, 2006)
95. A. Naumovich
On a finite volume discretization of the three-dimensional Biot poroelasticity system in multilayered domains
Keywords: Biot poroelasticity system, interface problems, finite volume discretization, finite difference method. (21 pages, 2006)
96. M. Krekel, J. Wenzel
A unified approach to Credit Default Swaption and Constant Maturity Credit Default Swap valuation
Keywords: LIBOR market model, credit risk, Credit Default Swaption, Constant Maturity Credit Default Swap-method. (43 pages, 2006)
97. A. Dreyer
Interval Methods for Analog Circuits
Keywords: interval arithmetic, analog circuits, tolerance analysis, parametric linear systems, frequency response, symbolic analysis, CAD, computer algebra (36 pages, 2006)
98. N. Weigel, S. Weihe, G. Bitsch, K. Dreßler
Usage of Simulation for Design and Optimization of Testing
Keywords: Vehicle test rigs, MBS, control, hydraulics, testing philosophy (14 pages, 2006)
99. H. Lang, G. Bitsch, K. Dreßler, M. Speckert
Comparison of the solutions of the elastic and elastoplastic boundary value problems
Keywords: Elastic BVP, elastoplastic BVP, variational inequalities, rate-independency, hysteresis, linear kinematic hardening, stop- and play-operator (21 pages, 2006)
100. M. Speckert, K. Dreßler, H. Mauch
MBS Simulation of a hexapod based suspension test rig
Keywords: Test rig, MBS simulation, suspension, hydraulics, controlling, design optimization (12 pages, 2006)
101. S. Azizi Sultan, K.-H. Küfer
A dynamic algorithm for beam orientations in multicriteria IMRT planning
Keywords: radiotherapy planning, beam orientation optimization, dynamic approach, evolutionary algorithm, global optimization (14 pages, 2006)
102. T. Götz, A. Klar, N. Marheineke, R. Wegener
A Stochastic Model for the Fiber Lay-down Process in the Nonwoven Production
Keywords: fiber dynamics, stochastic Hamiltonian system, stochastic averaging (17 pages, 2006)
103. Ph. Süß, K.-H. Küfer
Balancing control and simplicity: a variable aggregation method in intensity modulated radiation therapy planning
Keywords: IMRT planning, variable aggregation, clustering methods (22 pages, 2006)
104. A. Beaudry, G. Laporte, T. Melo, S. Nickel
Dynamic transportation of patients in hospitals
Keywords: in-house hospital transportation, dial-a-ride, dynamic mode, tabu search (37 pages, 2006)
105. Th. Hanne
Applying multiobjective evolutionary algorithms in industrial projects
Keywords: multiobjective evolutionary algorithms, discrete optimization, continuous optimization, electronic circuit design, semi-infinite programming, scheduling (18 pages, 2006)
106. J. Franke, S. Halim
Wild bootstrap tests for comparing signals and images
Keywords: wild bootstrap test, texture classification, textile quality control, defect detection, kernel estimate, nonparametric regression (13 pages, 2007)
107. Z. Drezner, S. Nickel
Solving the ordered one-median problem in the plane
Keywords: planar location, global optimization, ordered median, big triangle small triangle method, bounds, numerical experiments (21 pages, 2007)
108. Th. Götz, A. Klar, A. Unterreiter, R. Wegener
Numerical evidence for the non-existing of solutions of the equations describing rotational fiber spinning
Keywords: rotational fiber spinning, viscous fibers, boundary value problem, existence of solutions (11 pages, 2007)
109. Ph. Süß, K.-H. Küfer
Smooth intensity maps and the Bortfeld-Boyer sequencer
Keywords: probabilistic analysis, intensity modulated radiotherapy treatment (IMRT), IMRT plan application, step-and-shoot sequencing (8 pages, 2007)
110. E. Ivanov, O. Gluchshenko, H. Andrä, A. Kudryavtsev
Parallel software tool for decomposing and meshing of 3d structures
Keywords: a-priori domain decomposition, unstructured grid, Delaunay mesh generation (14 pages, 2007)
111. O. Iliev, R. Lazarov, J. Willems
Numerical study of two-grid preconditioners for 1d elliptic problems with highly oscillating discontinuous coefficients
Keywords: two-grid algorithm, oscillating coefficients, preconditioner (20 pages, 2007)
112. L. Bonilla, T. Götz, A. Klar, N. Marheineke, R. Wegener
Hydrodynamic limit of the Fokker-Planck equation describing fiber lay-down processes
Keywords: stochastic differential equations, Fokker-Planck equation, asymptotic expansion, Ornstein-Uhlenbeck process (17 pages, 2007)
113. S. Rief
Modeling and simulation of the pressing section of a paper machine
Keywords: paper machine, computational fluid dynamics, porous media (41 pages, 2007)
114. R. Ciegis, O. Iliev, Z. Lakdawala
On parallel numerical algorithms for simulating industrial filtration problems
Keywords: Navier-Stokes-Brinkmann equations, finite volume discretization method, SIMPLE, parallel computing, data decomposition method (24 pages, 2007)
115. N. Marheineke, R. Wegener
Dynamics of curved viscous fibers with surface tension
Keywords: Slender body theory, curved viscous fibers with surface tension, free boundary value problem (25 pages, 2007)

116. S. Feth, J. Franke, M. Speckert

**Resampling-Methoden zur mse-Korrektur
und Anwendungen in der Betriebsfestigkeit**

Keywords: Weibull, Bootstrap, Maximum-Likelihood,
Betriebsfestigkeit

(16 pages, 2007)

117. H. Knaf

**Kernel Fisher discriminant functions – a con-
cise and rigorous introduction**

Keywords: wild bootstrap test, texture classification,
textile quality control, defect detection, kernel estimate,
nonparametric regression

(30 pages, 2007)

Status quo: June 2007