



**Fraunhofer** Institut  
Techno- und  
Wirtschaftsmathematik

O. Iliev, I. Rybak

On approximation property of  
multipoint flux approximation  
method

© Fraunhofer-Institut für Techno- und Wirtschaftsmathematik ITWM 2007

ISSN 1434-9973

Bericht 119 (2007)

Alle Rechte vorbehalten. Ohne ausdrückliche schriftliche Genehmigung des Herausgebers ist es nicht gestattet, das Buch oder Teile daraus in irgendeiner Form durch Fotokopie, Mikrofilm oder andere Verfahren zu reproduzieren oder in eine für Maschinen, insbesondere Datenverarbeitungsanlagen, verwendbare Sprache zu übertragen. Dasselbe gilt für das Recht der öffentlichen Wiedergabe.

Warennamen werden ohne Gewährleistung der freien Verwendbarkeit benutzt.

Die Veröffentlichungen in der Berichtsreihe des Fraunhofer ITWM können bezogen werden über:

Fraunhofer-Institut für Techno- und  
Wirtschaftsmathematik ITWM  
Fraunhofer-Platz 1

67663 Kaiserslautern  
Germany

Telefon: +49 (0) 6 31/3 16 00-0  
Telefax: +49 (0) 6 31/3 16 00-10 99  
E-Mail: [info@itwm.fraunhofer.de](mailto:info@itwm.fraunhofer.de)  
Internet: [www.itwm.fraunhofer.de](http://www.itwm.fraunhofer.de)

# Vorwort

Das Tätigkeitsfeld des Fraunhofer-Instituts für Techno- und Wirtschaftsmathematik ITWM umfasst anwendungsnahe Grundlagenforschung, angewandte Forschung sowie Beratung und kundenspezifische Lösungen auf allen Gebieten, die für Techno- und Wirtschaftsmathematik bedeutsam sind.

In der Reihe »Berichte des Fraunhofer ITWM« soll die Arbeit des Instituts kontinuierlich einer interessierten Öffentlichkeit in Industrie, Wirtschaft und Wissenschaft vorgestellt werden. Durch die enge Verzahnung mit dem Fachbereich Mathematik der Universität Kaiserslautern sowie durch zahlreiche Kooperationen mit internationalen Institutionen und Hochschulen in den Bereichen Ausbildung und Forschung ist ein großes Potenzial für Forschungsberichte vorhanden. In die Berichtreihe sollen sowohl hervorragende Diplom- und Projektarbeiten und Dissertationen als auch Forschungsberichte der Institutsmitarbeiter und Institutsgäste zu aktuellen Fragen der Techno- und Wirtschaftsmathematik aufgenommen werden.

Darüber hinaus bietet die Reihe ein Forum für die Berichterstattung über die zahlreichen Kooperationsprojekte des Instituts mit Partnern aus Industrie und Wirtschaft.

Berichterstattung heißt hier Dokumentation des Transfers aktueller Ergebnisse aus mathematischer Forschungs- und Entwicklungsarbeit in industrielle Anwendungen und Softwareprodukte – und umgekehrt, denn Probleme der Praxis generieren neue interessante mathematische Fragestellungen.



Prof. Dr. Dieter Prätzel-Wolters  
Institutsleiter

Kaiserslautern, im Juni 2001



# ON APPROXIMATION PROPERTY OF MULTIPOINT FLUX APPROXIMATION METHOD

O.P. ILIEV AND I.V. RYBAK

ABSTRACT. Approximation property of multipoint flux approximation (MPFA) approach for elliptic equations with discontinuous full tensor coefficients is discussed here. Finite volume discretization of the above problem is presented in the case of jump discontinuities for the permeability tensor. First order approximation for the fluxes is proved. Results from numerical experiments are presented and discussed.

*Keywords:* Multipoint flux approximation, finite volume method, elliptic equation, discontinuous coefficients, anisotropy

## 1. INTRODUCTION

The paper concerns the approximation property of multipoint flux approximation (MPFA) approach applied to second order elliptic equation with discontinuous tensor coefficients. MPFA is a finite volume discretization in which the flux over a face is discretized using several grid points, in contrast to the standard two point approximation for the flux in the case of orthotropic problems. Recall that orthotropic problems are characterized by a diagonal coefficients tensor, while full tensor describes the coefficients of anisotropic problems. One of the first articles introducing MPFA were [8, 10], where MPFA was used in conjunction with simulation of porous media flow. No theoretical analysis was presented there. During the years, the method became popular due to its robustness and good accuracy. It was extensively used for solving applied problems, especially in geoscience. This risen the interest in a theoretical analysis of this approach, and some approximation and convergence results were published recently by Aavatsmark [1, 2], Klausen, Winther [4, 5], Wheeler, Yotov [3, 9]. Although the method was originally derived especially for problems with discontinuous tensor coefficients, the presented theoretical results concern MPFA applied to problems with continuous tensor coefficients. There are only numerical studies of the convergence order for discontinuous coefficients. The theoretical results presented in the above articles were obtained by showing equivalence of MPFA approximation to certain mixed finite element method, what allowed to use the well developed theory of mixed finite element method. First order convergence in  $W_2^1$  was proven, recall that the analysis was done for the case of smooth coefficients.

In the current article, we consider MPFA in the context of the theory of finite difference schemes [7, 11, 12]. We prove first order convergence for the MPFA method in discrete  $W_2^1$  norm. More precisely, we write down a specific derivation of MPFA, and show that the components of the continuous flux,  $\mathbf{W} = (W_1, W_2)^T$ , are approximated with  $O(h)$  in the midpoints of the edges by the components of the discrete flux,  $\mathbf{W}_h = (W_{1,h}, W_{2,h})^T$ . Further on, we use a priori estimates like (14) on

---

*Date:* November 27, 2007.

p.14 in [12]. This estimate allows to obtain convergence result in discrete  $W_2^1$  norm, if the error of a finite difference scheme can be written in divergence form, and if the approximation error for the fluxes is known. as it is often done in the analysis of finite difference schemes, we assume that the solution is piecewise smooth even when the coefficients have jump discontinuity. Obviously, our analysis is valid also for the case applying MPFA method for problems with continuous tensor coefficients. When the coefficients are discontinuous, often the solution has not enough regularity. However, for particular combination of the discontinuous coefficients, the solution may be smooth enough for our needs. We refer to [6] for a detailed discussion on the regularity of the solution in the case of scalar discontinuous coefficient.

So, we show that  $W_1(x_{i+1/2}, y_j) - W_{1,h}(x_{i+1/2}, y_j) = O(h)$ , where  $(x_{i+1/2}, y_j)$  is the middle point of an edge. We proceed with MPFA written in a specific way.

We consider a cell-centered basic grid, as well as a dual grid. The finite volume discretization is obtained on the original grid by applying divergence theorem. The flux over an edge is split into two parts, so that we have two subfluxes contributions over each edge: one for  $(x_{i+1/2}, y_j), (x_{i+1/2}, y_{j+1/2})$ , and another for  $(x_{i+1/2}, y_{j-1/2}), (x_{i+1/2}, y_j)$ . We will show the approximation only for the first subflux. Now we continue the discretization over the dual grid. For each of the four subcells of the dual cell, we will build **interpolating** linear polynomials. They interpolate the solution in the nodes of the original grid, and in the middle of the edges of the cells from the original grid, e.g., in  $(x_i, y_j), (x_{i+1/2}, y_j)$  and in  $(x_i, y_{j+1/2})$ . In fact, this is similar to one of the ways in which MPFA is derived by solving system of 4 equations with 4 unknowns. Our aim will be to show that the normal component of the flux,  $W^1$  in this case, is approximated with  $O(h)$  by the flux of this interpolating polynomial.

## 2. CONTINUOUS PROBLEM AND FINITE VOLUME DISCRETIZATION

In this section, we describe the mathematical model and the discretization approach. The derivation is based on the finite volume method (method of balance) and on multipoint flux approximation approach (MPFA).

**2.1. Statement of the problem.** In rectangular domain  $\bar{\Omega} = \Omega \cup \partial\Omega$ , we consider two-dimensional pressure equation obtained by combining the continuity equation ( $\nabla \cdot \mathbf{W} = f$ ) and Darcy's law ( $\mathbf{W} = -K\nabla u$ ) for steady state incompressible single phase flows in porous media

$$(1) \quad -\nabla \cdot (K\nabla u) = f, \quad \text{in } \Omega.$$

Boundary conditions complete the formulation of the problem

$$(2) \quad u = g^D, \quad \text{on } \Gamma_D, \quad K\nabla u \cdot \mathbf{n} = g^N, \quad \text{on } \Gamma_N, \quad \partial\Omega = \Gamma_D \cup \Gamma_N.$$

Here  $u$  is unknown pressure,  $f$  is the source, the set  $\Gamma_D$  is non-empty and had positive surface measure, the permeability tensor is full, symmetric, and uniformly positive definite in  $\Omega$ :

$$K = \begin{pmatrix} k_{11} & k_{12} \\ k_{12} & k_{22} \end{pmatrix} > 0, \quad k_{12} \neq 0.$$

The entries of the permeability tensor  $K$  may have jump discontinuities along certain interfaces that are parallel to the coordinate planes and along these interfaces

the perfect contact interface conditions are satisfied

$$[u] = 0, \quad [K\nabla u \cdot \mathbf{n}] = 0,$$

where  $[v] = v(\eta + 0) - v(\eta - 0)$  for the interface  $\eta$ .

We suppose that the following ellipticity conditions are satisfied as well

$$(3) \quad 0 < c_1 \sum_{\alpha=1}^2 \xi_\alpha^2 \leq \sum_{\alpha, \beta=1}^2 k_{\alpha\beta} \xi_\alpha \xi_\beta \leq c_2 \sum_{\alpha=1}^2 \xi_\alpha^2, \quad c_1, c_2 > 0,$$

where  $c_1, c_2 > 0$  are positive constants,  $\xi = (\xi_1, \xi_2)$  is an arbitrary nonzero vector  $|\xi| = \xi_1^2 + \xi_2^2 \neq 0$ .

*Remark.* Equation (1) describes also heat conductivity in composite materials, diffusion in heterogeneous media, stationary distribution of electric and magnetic field, etc.

**2.2. Finite volume discretization.** The domain  $\Omega$  is partitioned into blocks  $\Omega_{ij}$  so that the discontinuities of the permeability tensor  $K$  are aligned with cell boundaries. The centers of the cells  $\Omega_{ij}$  are denoted by  $(x_i, y_j)$  and the cell vertexes are the points  $(x_i \pm \frac{1}{2}h_1, y_j \pm \frac{1}{2}h_2)$ . The mesh that will be used to approximate the pressure will include all cell centers  $(x_i, y_j)$ . This mesh will be called *primary mesh*  $\omega_h = \{(x_i, y_j) : \Omega_{ij}\}$ . Similarly we shall use also the mesh of all cell vertexes, called often *dual mesh*. The velocities will be calculated at the points  $(x_i \pm \frac{1}{2}h_1, y_j)$  and  $(x_i, y_j \pm \frac{1}{2}h_2)$ .

The continuity equation  $(\nabla \cdot \mathbf{W} = f)$  is integrated over control volume  $\Omega_{ij}$  and making use of the divergence theorem, we obtain

$$(4) \quad \int_{\Omega_{ij}} \nabla \cdot \mathbf{W} d\mathbf{x} = \int_{\Omega_{ij}} f d\mathbf{x} \quad \Rightarrow \quad \int_{\partial\Omega_{ij}} \mathbf{W} \cdot \mathbf{n} ds = \int_{\Omega_{ij}} f d\mathbf{x}.$$

Replacing the velocity  $\mathbf{W}$  in (4) by certain approximation involving  $u$  by using the Darcy's relation  $\mathbf{W} = -K\nabla u$  we get a conservative method [7]. In this approximation we assume that the unknowns (or degrees of freedom) are the values of the pressure at the cell centers and then use these values to recover the velocity  $\mathbf{W}$ . According to the multipoint flux approximation this is done in the following manner. First we split each control volume

$$\Omega_{ij} = \left( x_i - \frac{1}{2}h_1, x_i + \frac{1}{2}h_1 \right) \times \left( y_j - \frac{1}{2}h_2, y_j + \frac{1}{2}h_2 \right)$$

into 4 subvolumes  $\Omega_{ij}^I = (x_i - \frac{1}{2}h_1, x_i) \times (y_j, y_j + \frac{1}{2}h_2)$ ,  $\Omega_{ij}^{II} = (x_i - \frac{1}{2}h_1, x_i) \times (y_j, y_j - \frac{1}{2}h_2)$ ,  $\Omega_{ij}^{III} = (x_i, x_i + \frac{1}{2}h_1) \times (y_j, y_j + \frac{1}{2}h_2)$ ,  $\Omega_{ij}^{IV} = (x_i + \frac{1}{2}h_1, x_i) \times (y_j, y_j - \frac{1}{2}h_2)$ .

We take the pressure to be a linear function on each subvolume  $\Omega_{ij}^k$  so that

$$(5) \quad u = \alpha^k x + \beta^k y + \gamma^k, \quad k = \overline{1, 4}.$$

The coefficients  $\alpha^k, \beta^k$  and  $\gamma^k$  in (5) determined by the following conditions:

(1c) the polynomials interpolate pressure values at the volume centers;

(2c) the continuity of the pressure at the centers of the faces of the volume  $\Omega_{ij}$  and the pressure data on faces that are part of  $\Gamma_D$ ;

(3c) the continuity of the normal component of the velocity  $\mathbf{v}$  at the centers of the faces of the volume  $\Omega_{ij}$  and the boundary data for the normal velocity on faces on  $\Gamma_N$ .




FIGURE 1. Control volume: inner and boundary cells

Conditions (1c)–(3c) correspond to O-method with surface midpoints as continuity points [1]. These conditions are applied on a cell from the dual grid, i.e. a cell centered at a vertex point from the dual grid (see interaction volume on Fig. 1). These cells are of three categories: cells corresponding to internal vertices, cells corresponding to boundary vertices, and 4 corner points of the domain  $\Omega$ .

Consider an internal vertex that is surrounded by four subcells with  $u_{i,j+1}$ ,  $u_{i+1,j+1}$ ,  $u_{i,j}$ ,  $u_{i+1,j}$  at the corners. To find the polynomial coefficients from (5), we use conditions (1c)–(3c). Note that four interaction volumes should be considered in order to find  $\alpha^k$ ,  $\beta^k$ ,  $\gamma^k$ ,  $k = \overline{1,4}$ .

The interpolating polynomials are defined by

$$\begin{aligned}
 P^1(x, y) &= \frac{u_{i+1/2,j} - u_{i,j}}{0.5h_1}(x + 0.5h_1) + \frac{u_{i,j+1/2} - u_{i,j}}{0.5h_2}(y + 0.5h_2) + u_{i,j} \\
 &= a^1x + b^1y + c^1, \\
 P^2(x, y) &= \frac{u_{i+1,j} - u_{i+1/2,j}}{0.5h_1}(x - 0.5h_1) + \frac{u_{i+1,j+1/2} - u_{i+1,j}}{0.5h_2}(y + 0.5h_2) \\
 &+ u_{i+1,j} = a^2x + b^2y + c^2, \\
 P^3(x, y) &= \frac{u_{i+1,j+1} - u_{i+1/2,j+1}}{0.5h_1}(x - 0.5h_1) + \frac{u_{i+1,j+1} - u_{i+1,j+1/2}}{0.5h_2}(y - 0.5h_2) \\
 &+ u_{i+1,j+1} = a^3x + b^3y + c^3, \\
 P^4(x, y) &= \frac{u_{i+1/2,j+1} - u_{i,j+1}}{0.5h_1}(x + 0.5h_1) + \frac{u_{i,j+1} - u_{i,j+1/2}}{0.5h_2}(y - 0.5h_2) \\
 &+ u_{i,j+1} = a^4x + b^4y + c^4,
 \end{aligned}
 \tag{6}$$

where  $u_{i+1/2,j} = u(x_{i+1/2}, y_j)$ ,  $u_{i,j} = u(x_i, y_j)$ , etc.

Similar to the condition for continuity of normal component of fluxes of the solution, we require these *interpolating* polynomials also to satisfy condition for continuity of normal component of their fluxes:

$$\begin{aligned}
 &k_{11} \frac{u_{i+1/2,j} - u_{i,j}}{0.5h_1} + k_{12} \frac{u_{i,j+1/2} - u_{i,j}}{0.5h_2} = \\
 &= k_{11}^{(+11)} \frac{u_{i+1,j} - u_{i+1/2,j}}{0.5h_1} + k_{12}^{(+11)} \frac{u_{i+1,j+1/2} - u_{i+1,j}}{0.5h_2}, \\
 &k_{12} \frac{u_{i+1/2,j} - u_{i,j}}{0.5h_1} + k_{22} \frac{u_{i,j+1/2} - u_{i,j}}{0.5h_2} = \\
 &= k_{12}^{(+12)} \frac{u_{i+1/2,j+1} - u_{i,j+1}}{0.5h_1} + k_{22}^{(+12)} \frac{u_{i,j+1} - u_{i,j+1/2}}{0.5h_2},
 \end{aligned}
 \tag{7}$$



$$\begin{aligned}
 & k_{11}^{(+12)} \frac{u_{i+1/2,j+1} - u_{i,j+1}}{0.5h_1} + k_{12}^{(+12)} \frac{u_{i,j+1} - u_{i,j+1/2}}{0.5h_2} = \\
 = & k_{11}^{(+11,+12)} \frac{u_{i+1,j+1} - u_{i+1/2,j+1}}{0.5h_1} + k_{12}^{(+11,+12)} \frac{u_{i+1,j+1} - u_{i+1,j+1/2}}{0.5h_2}, \\
 & k_{12}^{(+11)} \frac{u_{i+1,j} - u_{i+1/2,j}}{0.5h_1} + k_{22}^{(+11)} \frac{u_{i+1,j+1/2} - u_{i+1,j}}{0.5h_2} = \\
 = & k_{12}^{(+11,+12)} \frac{u_{i+1,j+1} - u_{i+1/2,j+1}}{0.5h_1} + k_{22}^{(+11,+12)} \frac{u_{i+1,j+1} - u_{i+1,j+1/2}}{0.5h_2}.
 \end{aligned}$$

As an auxiliary step, we write the above equalities as a system (7) with respect to unknown values at the midpoint of edges:

$$(8) \quad Av = w,$$

where  $v = (u_{i+1/2,j}, u_{i,j+1/2}, u_{i+1/2,j+1}, u_{i+1,j+1/2})^T$ ,  $w = (w_1, w_2, w_3, w_4)^T$ ,

$$A = \begin{pmatrix} \frac{k_{11} + k_{11}^{(+11)}}{h_1} & \frac{k_{12}}{h_2} & 0 & -\frac{k_{12}^{(+11)}}{h_2} \\ \frac{k_{12}}{h_1} & \frac{k_{22} + k_{22}^{(+12)}}{h_2} & -\frac{k_{12}^{(+12)}}{h_1} & 0 \\ 0 & -\frac{k_{12}^{(+12)}}{h_2} & \frac{k_{11}^{(+12)} + k_{11}^{(+11,+12)}}{h_1} & \frac{k_{12}^{(+11,+12)}}{h_2} \\ -\frac{k_{12}^{(+11)}}{h_1} & 0 & \frac{k_{12}^{(+11,+12)}}{h_1} & \frac{k_{22}^{(+11)} + k_{22}^{(+11,+12)}}{h_2} \end{pmatrix}.$$

Here

$$\begin{aligned}
 w_1 &= \left( \frac{k_{11}^{(+11)}}{h_1} - \frac{k_{12}^{(+11)}}{h_2} \right) u_{i+1,j} + \left( \frac{k_{11}}{h_1} + \frac{k_{12}}{h_2} \right) u_{i,j}, \\
 w_2 &= \left( \frac{k_{12}}{h_1} + \frac{k_{22}}{h_2} \right) u_{i,j} - \left( \frac{k_{12}^{(+12)}}{h_1} - \frac{k_{22}^{(+12)}}{h_2} \right) u_{i,j+1}, \\
 w_3 &= \left( \frac{k_{11}^{(+12)}}{h_1} - \frac{k_{12}^{(+12)}}{h_2} \right) u_{i,j+1} + \left( \frac{k_{11}^{(+11,+12)}}{h_1} + \frac{k_{12}^{(+11,+12)}}{h_2} \right) u_{i+1,j+1}, \\
 w_4 &= \left( -\frac{k_{12}^{(+11)}}{h_1} + \frac{k_{22}^{(+11)}}{h_2} \right) u_{i+1,j} + \left( \frac{k_{12}^{(+11,+12)}}{h_1} + \frac{k_{22}^{(+11,+12)}}{h_2} \right) u_{i+1,j+1}.
 \end{aligned}$$

Now, from the auxiliary step we come to reformulating the system in the form we need. The quantities which are of interest for us at this stage are the approximations to the derivatives of the solution. These new variables coincide with the coefficients of the polynomials  $P^i(x, y)$ :

$$\begin{aligned}
 a^1 &= \frac{u_{i+1/2,j} - u_{i,j}}{0.5h_1}, & b^1 &= \frac{u_{i,j+1/2} - u_{i,j}}{0.5h_2}, \\
 a^3 &= \frac{u_{i+1,j+1} - u_{i+1/2,j+1}}{0.5h_1}, & b^3 &= \frac{u_{i+1,j+1} - u_{i+1,j+1/2}}{0.5h_2}.
 \end{aligned}$$

So, we rewrite system (8) in new variables  $\tilde{A}\tilde{v} = \tilde{w}$  with  $\tilde{v} = (a^1, b^1, a^3, b^3)^T$ . The matrix of the system is

$$(9) \quad \tilde{A} = \begin{pmatrix} k_{11} + k_{11}^{(+11)} & k_{12} & 0 & k_{12}^{(+11)} \\ k_{12} & k_{22} + k_{22}^{(+12)} & k_{12}^{(+12)} & 0 \\ 0 & k_{12}^{(+12)} & k_{11}^{(+12)} + k_{11}^{(+11,+12)} & k_{12}^{(+11,+12)} \\ k_{12}^{(+11)} & 0 & k_{12}^{(+11,+12)} & k_{22}^{(+11)} + k_{22}^{(+11,+12)} \end{pmatrix},$$

and the right-hand side is given by

$$\begin{aligned}\tilde{w}_1 &= 2 \left( k_{11}^{(+11)} \frac{u_{i+1,j} - u_{i,j}}{h_1} + k_{12}^{(+11)} \frac{u_{i+1,j+1} - u_{i+1,j}}{h_2} \right), \\ \tilde{w}_2 &= 2 \left( k_{yx}^{(+12)} \frac{u_{i+1,j+1} - u_{i,j+1}}{h_1} + k_{22}^{(+12)} \frac{u_{i,j+1} - u_{i,j}}{h_2} \right), \\ \tilde{w}_3 &= 2 \left( k_{11}^{(+12)} \frac{u_{i+1,j+1} - u_{i,j+1}}{h_1} + k_{12}^{(+12)} \frac{u_{i,j+1} - u_{i,j}}{h_2} \right), \\ \tilde{w}_4 &= 2 \left( k_{12}^{(+11)} \frac{u_{i+1,j} - u_{i,j}}{h_1} + k_{22}^{(+11)} \frac{u_{i+1,j+1} - u_{i+1,j}}{h_2} \right).\end{aligned}$$

Coefficients  $a^i$ ,  $b^i$ ,  $i = 2, 4$ , can be expressed through  $a^i$ ,  $b^i$ ,  $i = 1, 3$  and the pressure values at the cell centers

$$(10) \quad \begin{aligned}a^2 &= \frac{u_{i+1,j} - u_{i,j}}{0.5h_1} - a^1, & b^2 &= \frac{u_{i+1,j+1} - u_{i+1,j}}{0.5h_2} - b^3 \\ a^4 &= \frac{u_{i+1,j+1} - u_{i,j+1}}{0.5h_1} - a^3, & b^4 &= \frac{u_{i,j+1} - u_{i,j}}{0.5h_2} - b^1.\end{aligned}$$

Note, that it gives us the expressions for the velocity that is constant over each of the 4 subcells of the vertex-centered volume. Consider in the same way three other vertex-centered volumes to find the fluxes incoming and outgoing the cell-centered (control volume)  $\Omega_{ij}$ . These formulas are used to find  $\mathbf{W} \cdot \mathbf{n}$  on  $\partial\Omega_{ij}$ , as needed by the relation (4).

We use the midpoint rule for calculating the integrals in the balance method (4), that provides

$$h_2 W_{1,h}^{out} - h_2 W_{1,h}^{in} + h_1 W_{2,h}^{out} - h_1 W_{2,h}^{in} = h_1 h_2 f.$$

Here  $W_{i,h}^{in}$ ,  $W_{i,h}^{out}$ ,  $i = 1, 2$  are obtained by summing the two fluxes incoming and leaving the considered control volume (Fig. 1). Thus, the difference scheme for the pressure equation (1) can be written in the following form

$$(11) \quad \begin{aligned}& \frac{h_2}{2} \left( k_{11} a^2 + k_{12} b^2 + k_{11}^{(+11)} \check{a}^3 + k_{12}^{(+11)} \check{b}^3 \right) - \\ & \frac{h_2}{2} \left( k_{11}^{(-11)} \bar{a}^2 + k_{12}^{(-11)} \bar{b}^2 + k_{11} \check{\bar{a}}^3 + k_{12} \check{\bar{b}}^3 \right) + \\ & \frac{h_1}{2} \left( k_{12} a^2 + k_{22} b^2 + k_{12}^{(+12)} \bar{a}^3 + k_{22}^{(+12)} \bar{b}^3 \right) - \\ & \frac{h_2}{2} \left( k_{12}^{(-12)} \check{\bar{a}}^2 - k_{22}^{(-12)} \check{\bar{b}}^2 - k_{12} \check{\bar{a}}^3 - k_{22} \check{\bar{b}}^3 \right) = -f h_1 h_2,\end{aligned}$$

where coefficients  $\bar{a}^i$ ,  $\bar{b}^i$  are calculated by the same formulas as the coefficients  $a^i$ ,  $b^i$ , but in the grid block  $u_{i-1,j+1}u_{i,j+1}u_{i,j}u_{i-1,j}$ . The coefficients  $\check{a}^i$ ,  $\check{b}^i$  and  $\check{\bar{a}}^i$ ,  $\check{\bar{b}}^i$  are calculated in the grid blocks  $u_{i,j}u_{i+1,j}u_{i+1,j-1}u_{i,j-1}$  and  $u_{i-1,j}u_{i,j}u_{i,j-1}u_{i-1,j-1}$  respectively.

For vertex that is on the boundary the situation is simpler. In the case of Neumann boundary conditions the flux is given on the boundary, while in the case of Dirichlet boundary conditions we just do the same procedure as for the inner control volume, but in this case we have only one interface.

Combining this relationship for each neighboring vertex gives us a discrete pressure equation with a 9-point stencil.

*Remark.* In the case of diagonal (orthotropic) permeability tensor ( $k_{12} = 0$ ), scheme (11) reduces to the well-known harmonic averaging finite difference scheme

$$\begin{aligned} & \frac{1}{h_1} \left( \frac{2k_{11}k_{11}^{(+11)}}{k_{11} + k_{11}^{(+11)}} \frac{u_{i+1,j} - u_{i,j}}{h_1} - \frac{2k_{11}k_{11}^{(-11)}}{k_{11} + k_{11}^{(-11)}} \frac{u_{i,j} - u_{i-1,j}}{h_1} \right) + \\ & \frac{1}{h_2} \left( \frac{2k_{22}k_{22}^{(+12)}}{k_{22} + k_{22}^{(+12)}} \frac{u_{i,j+1} - u_{i,j}}{h_2} - \frac{2k_{22}k_{22}^{(-12)}}{k_{22} + k_{22}^{(-12)}} \frac{u_{i,j} - u_{i,j-1}}{h_2} \right) = -f. \end{aligned}$$

*Remark.* In the case, the coefficients  $k_{11}$ ,  $k_{12}$ ,  $k_{22}$  are constant, the difference scheme (11) can be written in the following form

$$k_{11}u_{x\bar{x}} + k_{12}\frac{u_{\bar{x}y} + u_{x\bar{y}}}{2} + k_{12}\frac{u_{\bar{x}\bar{y}} + u_{xy}}{2} + k_{xy}u_{y\bar{y}} + R = -f,$$

where

$$R = \frac{k_{12}^2}{4} \left( \frac{h_1^2}{k_{11}} + \frac{h_2^2}{k_{22}} \right) u_{\bar{x}\bar{x}\bar{y}y} = O(h^2).$$

So, in this case the finite volume scheme (11) is equivalent to the second-order finite difference scheme from [7]:

$$k_{11}u_{x\bar{x}} + 0.5k_{12}(u_{\bar{x}y} + u_{x\bar{y}} + u_{\bar{x}\bar{y}} + u_{xy}) + k_{22}u_{y\bar{y}} = -f.$$

### 3. APPROXIMATION AND CONVERGENCE PROPERTIES

In this section, we prove the first order approximation for the fluxes in the middle points of the edges in the case of discontinuous tensor coefficients.

**3.1. Approximation order of fluxes.** We examine the order of the approximation of the fluxes in the case of discontinuous tensor coefficients and piecewise smooth solution. To do this, consider the condition for the continuity of normal component of fluxes of the PDE solution at midpoints of edges, and its discrete approximation be the normal components of fluxes of the above derived piecewise linear polynomials. By considering the difference of the both, we will get expressions for the approximation error. After certain manipulations, we show that the approximation is  $O(h)$ .

To start, consider the flux continuity condition at the point  $(x_{i+1/2}, y_j)$ :

$$(12) \quad \begin{aligned} & k_{11} \frac{\partial u}{\partial x} \Big|_{(x_{i+1/2-0}, y_j)} + k_{12} \frac{\partial u}{\partial y} \Big|_{(x_{i+1/2-0}, y_j)} = \\ & k_{11}^{(+11)} \frac{\partial u}{\partial x} \Big|_{(x_{i+1/2+0}, y_j)} + k_{12}^{(+11)} \frac{\partial u}{\partial y} \Big|_{(x_{i+1/2+0}, y_j)}, \end{aligned}$$

and its discrete approximation (see formula (7)):

$$(13) \quad \begin{aligned} & k_{11} \frac{u_{i+1/2,j} - u_{i,j}}{0.5h_1} + k_{12} \frac{u_{i,j+1/2} - u_{i,j}}{0.5h_2} = \\ & k_{11}^{(+11)} \frac{u_{i+1,j} - u_{i+1/2,j}}{0.5h_1} + k_{12}^{(+11)} \frac{u_{i+1,j+1/2} - u_{i+1,j}}{0.5h_2}. \end{aligned}$$

By subtracting equation (12) from (13), we obtain

$$(14) \quad \begin{aligned} & k_{11} \left( \frac{u_{i+1/2,j} - u_{i,j}}{0.5h_1} - \frac{\partial u}{\partial x} \Big|_{(x_{i+1/2-0}, y_j)} \right) + \\ & k_{12} \left( \frac{u_{i,j+1/2} - u_{i,j}}{0.5h_2} - \frac{\partial u}{\partial y} \Big|_{(x_{i+1/2-0}, y_j)} \right) = \\ & k_{11}^{(+11)} \left( \frac{u_{i+1,j} - u_{i+1/2,j}}{0.5h_1} - \frac{\partial u}{\partial x} \Big|_{(x_{i+1/2+0}, y_j)} \right) + \\ & k_{12}^{(+11)} \left( \frac{u_{i+1,j+1/2} - u_{i+1,j}}{0.5h_2} - \frac{\partial u}{\partial y} \Big|_{(x_{i+1/2+0}, y_j)} \right). \end{aligned}$$

From polynomials (6) we have

$$\begin{aligned} \frac{u_{i+1/2,j} - u_{i,j}}{0.5h_1} &= \frac{\partial P^1}{\partial x}, & \frac{u_{i+1,j} - u_{i+1/2,j}}{0.5h_1} &= \frac{\partial P^2}{\partial x}, \\ \frac{u_{i,j+1/2} - u_{i,j}}{0.5h_2} &= \frac{\partial P^1}{\partial y}, & \frac{u_{i+1,j+1/2} - u_{i+1,j}}{0.5h_2} &= \frac{\partial P^2}{\partial y}. \end{aligned}$$

Substituting these expressions into equation (14), we get

$$(15) \quad \begin{aligned} & k_{11} \left( \frac{\partial P^1}{\partial x} - \frac{\partial u}{\partial x} \Big|_{(x_{i+1/2-0}, y_j)} \right) + k_{12} \left( \frac{\partial P^1}{\partial y} - \frac{\partial u}{\partial y} \Big|_{(x_{i+1/2-0}, y_j)} \right) = \\ & k_{11}^{(+11)} \left( \frac{\partial P^2}{\partial x} - \frac{\partial u}{\partial x} \Big|_{(x_{i+1/2+0}, y_j)} \right) + k_{12}^{(+11)} \left( \frac{\partial P^2}{\partial y} - \frac{\partial u}{\partial y} \Big|_{(x_{i+1/2+0}, y_j)} \right). \end{aligned}$$

Simple calculations give

$$\frac{\partial P^2}{\partial x} = \frac{u_{i+1,j} - u_{i,j}}{0.5h_1} - \frac{\partial P^1}{\partial x}.$$

Using the Taylor expansion at the point  $e$ , from the last expression we obtain

$$(16) \quad \frac{\partial P^2}{\partial x} = \frac{\partial u}{\partial x} \Big|_{(x_{i+1/2+0}, y_j)} + \frac{\partial u}{\partial x} \Big|_{(x_{i+1/2-0}, y_j)} - \frac{\partial P^1}{\partial x} + O(h_1).$$

Substituting the above relation in (15), we get

$$(17) \quad \begin{aligned} & k_{11} \left( \frac{\partial P^1}{\partial x} - \frac{\partial u}{\partial x} \Big|_{(x_{i+1/2-0}, y_j)} \right) + k_{12} \left( \frac{\partial P^1}{\partial y} - \frac{\partial u}{\partial y} \Big|_{(x_{i+1/2-0}, y_j)} \right) = \\ & k_{11}^{(+11)} \left( \left( \frac{\partial u}{\partial x} \Big|_{(x_{i+1/2+0}, y_j)} + \frac{\partial u}{\partial x} \Big|_{(x_{i+1/2-0}, y_j)} - \frac{\partial P^1}{\partial x} + O(h_1) \right) - \frac{\partial u}{\partial x} \Big|_{(x_{i+1/2+0}, y_j)} \right) + \\ & k_{12}^{(+11)} \left( \frac{\partial P^2}{\partial y} - \frac{\partial u}{\partial y} \Big|_{(x_{i+1/2+0}, y_j)} \right). \end{aligned}$$

Consider new variables,  $z_1, z_2$ , which are the approximation error for  $x$ -derivative at the point  $(x_{i+1/2-0}, y_j)$ , and for  $y$ -derivative of the solution at point  $(x_i, y_{j+1/2-0})$ , respectively:

$$z_1 = \left( \frac{\partial P^1}{\partial x} - \frac{\partial u}{\partial x} \right) \Big|_{(x_{i+1/2-0}, y_j)}, \quad z_2 = \left( \frac{\partial P^1}{\partial y} - \frac{\partial u}{\partial y} \right) \Big|_{(x_i, y_{j+1/2-0})}.$$

Now, we assume that the solution is piecewise smooth within the cells, where the coefficients are piecewise constants. Under this assumption, we have

$$\left(\frac{\partial u}{\partial y}\right)\Big|_{(x_{i+1/2-0}, y_j)} = \left(\frac{\partial u}{\partial y}\right)\Big|_{(x_i, y_{j+1/2-0})} + O(h)$$

and respectively,

$$\left(\frac{\partial P^1}{\partial y} - \frac{\partial u}{\partial y}\right)\Big|_{(x_{i+1/2-0}, y_j)} = \left(\frac{\partial P^1}{\partial y} - \frac{\partial u}{\partial y}\right)\Big|_{(x_i, y_{j+1/2-0})} + O(h) = z_2 + O(h)$$

Thus, equation (17) can be rewritten as

$$(18) \quad \left(k_{11} + k_{11}^{(+11)}\right) z_1 + k_{12} z_2 = O(h_1) + k_{12}^{(+11)} \left(\frac{\partial P^2}{\partial y} - \frac{\partial u}{\partial y}\Big|_{(x_{i+1/2+0}, y_j)}\right).$$

We leave for a while the above equation in this unfinished form, and continue with relations at the point  $(x_i, y_{j+1/2})$ . Similarly to the above derivations for the point  $(x_{i+1/2}, y_j)$ , we consider the flux continuity condition at the point  $(x_i, y_{j+1/2})$ :

$$(19) \quad \begin{aligned} & k_{12} \left(\frac{u_{i+1/2, j} - u_{i, j}}{0.5h_1} - \frac{\partial u}{\partial x}\Big|_{(x_i, y_{j+1/2-0})}\right) + \\ & k_{22} \left(\frac{u_{i, j+1/2} - u_{i, j}}{0.5h_2} - \frac{\partial u}{\partial y}\Big|_{(x_i, y_{j+1/2-0})}\right) = \\ & k_{12}^{(+12)} \left(\frac{u_{i+1/2, j+1} - u_{i, j+1}}{0.5h_1} - \frac{\partial u}{\partial x}\Big|_{(x_i, y_{j+1/2+0})}\right) + \\ & k_{22}^{(+12)} \left(\frac{u_{i, j+1} - u_{i, j+1/2}}{0.5h_2} - \frac{\partial u}{\partial y}\Big|_{(x_i, y_{j+1/2+0})}\right). \end{aligned}$$

Recall that derivatives of the polynomial and  $P^4$  (6) are given by

$$\frac{u_{i+1/2, j+1} - u_{i, j+1}}{0.5h_1} = \frac{\partial P^4}{\partial x}, \quad \frac{u_{i, j+1} - u_{i, j+1/2}}{0.5h_2} = \frac{\partial P^4}{\partial y}.$$

Substituting these expressions together with similar expressions for  $P^1$  into equation (19), we get

$$(20) \quad \begin{aligned} & k_{12} \left(\frac{\partial P^1}{\partial x} - \frac{\partial u}{\partial x}\Big|_{(x_i, y_{j+1/2-0})}\right) + k_{22} \left(\frac{\partial P^1}{\partial y} - \frac{\partial u}{\partial y}\Big|_{(x_i, y_{j+1/2-0})}\right) = \\ & k_{12}^{(+12)} \left(\frac{\partial P^4}{\partial x} - \frac{\partial u}{\partial x}\Big|_{(x_i, y_{j+1/2+0})}\right) + k_{22}^{(+12)} \left(\frac{\partial P^4}{\partial y} - \frac{\partial u}{\partial y}\Big|_{(x_i, y_{j+1/2+0})}\right). \end{aligned}$$

It is easy to see that

$$\frac{\partial P^4}{\partial y} = \frac{u_{i, j+1} - u_{i, j}}{0.5h_2} - \frac{\partial P^1}{\partial y} = \frac{\partial u}{\partial y}\Big|_{(x_i, y_{j+1/2+0})} + \frac{\partial u}{\partial y}\Big|_{(x_i, y_{j+1/2-0})} - \frac{\partial P^1}{\partial y} + O(h_2).$$

Substituting this into (20), and using the expansion

$$\left(\frac{\partial P^1}{\partial x} - \frac{\partial u}{\partial x}\right)\Big|_{(x_i, y_{j+1/2-0})} = \left(\frac{\partial P^1}{\partial x} - \frac{\partial u}{\partial x}\right)\Big|_{(x_{i+1/2-0}, y_j)} + O(h) = z_1 + O(h),$$

we get

$$(21) \quad k_{12} z_1 + \left(k_{22} + k_{22}^{(+12)}\right) z_2 = k_{12}^{(+12)} \left(\frac{\partial P^4}{\partial x} - \frac{\partial u}{\partial x}\Big|_{(x_i, y_{j+1/2+0})}\right) + O(h).$$

Now we proceed with the remaining two edges,  $(x_{i+1/2}, y_{j+1})$  and  $(x_{i+1}, y_{j+1/2-0})$ , as well as with remaining terms in the equations at  $(x_{i+1/2+0}, y_j)$  and  $(x_i, y_{j+1/2+0})$ . First, we introduce the variables

$$z_3 = \left( \frac{\partial P^3}{\partial x} - \frac{\partial u}{\partial x} \right) \Big|_{(x_{i+1/2+0}, y_{j+1})}, \quad z_4 = \left( \frac{\partial P^3}{\partial y} - \frac{\partial u}{\partial y} \right) \Big|_{(x_{i+1}, y_{j+1/2+0})},$$

By simple calculations we get

$$(22) \quad \frac{\partial P^2}{\partial y} = \frac{\partial u}{\partial y} \Big|_{(x_{i+1}, y_{j+1/2+0})} + \frac{\partial u}{\partial y} \Big|_{(x_{i+1}, y_{j+1/2-0})} - \frac{\partial P^3}{\partial y} + O(h_2),$$

$$(23) \quad \frac{\partial P^4}{\partial x} = \frac{\partial u}{\partial x} \Big|_{(x_{i+1/2+0}, y_{j+1})} + \frac{\partial u}{\partial x} \Big|_{(x_{i+1/2-0}, y_{j+1})} - \frac{\partial P^3}{\partial x} + O(h_1).$$

Next, from the assumption that the solution is piecewise smooth, we have

$$\frac{\partial u}{\partial x} \Big|_{(x_{i+1}, y_{j+1/2-0})} = \frac{\partial u}{\partial x} \Big|_{(x_{i+1/2+0}, y_j)} + O(h),$$

$$\frac{\partial u}{\partial y} \Big|_{(x_{i+1/2+0}, y_j)} = \frac{\partial u}{\partial y} \Big|_{(x_{i+1}, y_{j+1/2-0})} + O(h).$$

Then using the above formula together with formula (22), we obtain

$$\begin{aligned} \frac{\partial P^2}{\partial y} - \frac{\partial u}{\partial y} \Big|_{(x_{i+1/2+0}, y_j)} &= \frac{\partial P^2}{\partial y} - \frac{\partial u}{\partial y} \Big|_{(x_{i+1}, y_{j+1/2-0})} + O(h) = \\ &= - \left( \frac{\partial P^3}{\partial y} - \frac{\partial u}{\partial y} \Big|_{(x_{i+1}, y_{j+1/2+0})} \right) + O(h) = -z_4 + O(h). \end{aligned}$$

Thus, equation (18) can be written as

$$\left( k_{11} + k_{11}^{(+11)} \right) z_1 + k_{12} z_2 + k_{12}^{(+11)} z_4 = O(h_1).$$

In the same way we obtain

$$\begin{aligned} \frac{\partial P^4}{\partial x} - \frac{\partial u}{\partial x} \Big|_{(x_i, y_{j+1/2+0})} &= \frac{\partial P^4}{\partial x} - \frac{\partial u}{\partial x} \Big|_{(x_{i+1/2-0}, y_{j+1})} + O(h) = \\ &= - \left( \frac{\partial P^3}{\partial x} - \frac{\partial u}{\partial x} \Big|_{(x_{i+1/2+0}, y_{j+1})} \right) + O(h) = -z_3 + O(h), \end{aligned}$$

and rewrite equation (21) in the following way

$$k_{12} z_1 + \left( k_{22} + k_{22}^{(+12)} \right) z_2 + k_{12}^{(+12)} z_3 = O(h).$$

The last what we need before finalizing the derivations, are the continuity conditions at the points  $(x_{i+1/2}, y_{j+1})$  and  $(x_{i+1}, y_{j+1/2-0})$ :

$$(24) \quad \begin{aligned} &k_{11}^{(+12)} \left( \frac{\partial P^4}{\partial x} - \frac{\partial u}{\partial x} \Big|_{(x_{i+1/2-0}, y_{j+1})} \right) + k_{12}^{(+12)} \left( \frac{\partial P^4}{\partial y} - \frac{\partial u}{\partial y} \Big|_{(x_i, y_{j+1/2+0})} \right) = \\ &k_{11}^{(+11, +12)} \left( \frac{\partial P^3}{\partial x} - \frac{\partial u}{\partial x} \Big|_{(x_{i+1/2+0}, y_{j+1})} \right) + k_{12}^{(+11, +12)} \left( \frac{\partial P^3}{\partial y} - \frac{\partial u}{\partial y} \Big|_{(x_{i+1}, y_{j+1/2+0})} \right). \end{aligned}$$

$$(25) \quad \begin{aligned} & k_{12}^{(+11)} \left( \frac{\partial P^2}{\partial x} - \frac{\partial u}{\partial x} \Big|_{(x_{i+1/2+0}, y_j)} \right) + k_{22}^{(+11)} \left( \frac{\partial P^2}{\partial y} - \frac{\partial u}{\partial y} \Big|_{(x_{i+1}, y_{j+1/2-0})} \right) = \\ & k_{12}^{(+11,+12)} \left( \frac{\partial P^3}{\partial x} - \frac{\partial u}{\partial x} \Big|_{(x_{i+1/2+0}, y_{j+1})} \right) + k_{12}^{(+11,+12)} \left( \frac{\partial P^3}{\partial y} - \frac{\partial u}{\partial y} \Big|_{(x_{i+1}, y_{j+1/2+0})} \right). \end{aligned}$$

Using similar approach and taking into account that

$$\begin{aligned} \frac{u_{i+1,j+1} - u_{i+1/2,j+1}}{0.5h_1} &= \frac{\partial P^3}{\partial x}, \quad \frac{u_{i+1,j+1} - u_{i+1,j+1/2}}{0.5h_2} = \frac{\partial P^3}{\partial y}, \\ \frac{\partial u}{\partial x} \Big|_{(x_{i+1}, y_{j+1/2-0})} &= \frac{\partial u}{\partial x} \Big|_{(x_{i+1/2+0}, y_j)} + O(h), \quad \frac{\partial u}{\partial y} \Big|_{(x_{i+1/2-0}, y_{j+1})} = \frac{\partial u}{\partial y} \Big|_{(x_i, y_{j+1/2+0})} + O(h), \\ \left( \frac{\partial P^3}{\partial x} - \frac{\partial u}{\partial x} \right) \Big|_{(x_{i+1}, y_{j+1/2+0})} &= \left( \frac{\partial P^3}{\partial x} - \frac{\partial u}{\partial x} \right) \Big|_{(x_{i+1/2+0}, y_{j+1})} + O(h) = z_3 + O(h), \\ \left( \frac{\partial P^3}{\partial y} - \frac{\partial u}{\partial y} \right) \Big|_{(x_{i+1/2+0}, y_{j+1})} &= \left( \frac{\partial P^3}{\partial y} - \frac{\partial u}{\partial y} \right) \Big|_{(x_{i+1}, y_{j+1/2+0})} + O(h) = q_3 + O(h), \end{aligned}$$

we rewrite equations (24), (25) in the following form

$$\begin{aligned} k_{12}^{(+12)} z_1 + \left( k_{11}^{(+12)} + k_{11}^{(+11,+12)} \right) z_3 + k_{12}^{(+11,+12)} z_4 &= O(h), \\ k_{12}^{(+11)} z_1 + k_{12}^{(+11,+12)} z_3 + \left( k_{22}^{(+11)} + k_{22}^{(+11,+12)} \right) z_4 &= O(h). \end{aligned}$$

So, system (15), (20), (24), (25) can be written as

$$Mz = r,$$

where  $M = \tilde{A}$ ,  $z = (z_1, z_2, z_3, z_4)^T$ ,  $r = (r_1, r_2, r_3, r_4)^T$ , and  $r_i = O(h)$ . It is easy to see that matrix  $M$  is symmetric. To show that it is positive definite, consider the scalar product

$$(26) \quad \begin{aligned} (Mv, v) &= k_{11}^{(+11)} v_1^2 + k_{11} v_1^2 + 2k_{12} v_1 v_2 + k_{22}^{(+12)} v_2^2 + k_{22} v_2^2 + \\ & 2k_{12}^{(+12)} v_2 v_3 + k_{11}^{(+12)} v_3^2 + k_{11}^{(+11,+12)} v_3^2 + 2k_{12}^{(+11)} v_1 v_4 + \\ & 2k_{12}^{(+11,+12)} v_3 v_4 + k_{22}^{(+11)} v_4^2 + k_{22}^{(+11,+12)} v_4^2, \end{aligned}$$

where  $v = (v_1, v_2, v_3, v_4)^T$ . Since  $K$  is positive definite in every point, than taking into account the ellipticity condition (3) valid for any nonzero  $\xi = (\xi_1, \xi_2)$ ,  $|\xi| = \xi_1^2 + \xi_2^2 \neq 0$ , and using  $\xi = (v_1, v_2)$ ,  $\xi = (v_1, v_4)$ ,  $\xi = (v_3, v_2)$ ,  $\xi = (v_3, v_4)$ , from (26) we get

$$\begin{aligned} (Mv, v) &= (k_{11} v_1^2 + 2k_{12} v_1 v_2 + k_{22} v_2^2) + \left( k_{11}^{(+11)} v_1^2 + 2k_{12}^{(+11)} v_1 v_4 + k_{22}^{(+11)} v_4^2 \right) + \\ & \left( k_{11}^{(+12)} v_3^2 + 2k_{12}^{(+12)} v_2 v_3 + k_{22}^{(+12)} v_2^2 \right) + \\ & \left( k_{11}^{(+11,+12)} v_3^2 + 2k_{12}^{(+11,+12)} v_3 v_4 + k_{22}^{(+11,+12)} v_4^2 \right) \geq \\ & c_1 (v_1^2 + v_2^2) + c_1 (v_1^2 + v_4^2) + c_1 (v_3^2 + v_2^2) + c_1 (v_3^2 + v_4^2) = \\ & 2c_1 (v_1^2 + v_2^2 + v_3^2 + v_4^2) > 0. \end{aligned}$$

Hence, there exists  $M^{-1}$  such that  $z = \tilde{r}$ , where  $\tilde{r} = M^{-1}r$ , and  $\tilde{r}_i = O(h)$ . Thus

$$z_i = O(h), \quad q_i = O(h), \quad i = \overline{1,4},$$

and we have first order approximation for the normal components of the fluxes at the midpoints of the edges.

**3.2. Convergence of MPFA.** In this section we prove convergence of MPFA knowing that the flux is approximated with  $O(h)$ .

In the domain  $\bar{\Omega}$  we consider staggered mesh  $\bar{\omega}_0 = \omega_0 \cup \gamma_0$ ,  $\bar{\omega}_1 = \omega_1 \cup \gamma_1$ ,  $\bar{\omega}_2 = \omega_2 \cup \gamma_2$ . Here  $\omega_0$  defines the cell centers where the pressure is considered, while  $\omega_1$  and  $\omega_2$  define the midpoints of the edges, where the velocity components (i.e., fluxes) are defined.

$$\bar{\omega}_0 = \{(i + 1/2)h_1, (j + 1/2)h_2), i = -1/2, 0, 1, \dots, N_1, N_1 + 1/2,$$

$$j = -1/2, 0, 1, \dots, N_2, N_2 + 1/2\},$$

$$\bar{\omega}_1 = \{(ih_1, (j + 1/2)h_2), i = 0, 1, \dots, N_1, j = 0, 1, \dots, N_2\},$$

$$\bar{\omega}_2 = \{((i + 1/2)h_1, jh_2), i = 0, 1, \dots, N_1, j = 0, 1, \dots, N_2\},$$

where  $h_1 = 1/N_1$ ,  $h_2 = 1/N_2$ , and  $N_1, N_2$  are positive integers. Note that on the boundary there are nodes belonging to  $\omega_0$  and  $\omega_1$ , or  $\omega_0$  and  $\omega_2$ .




FIGURE 2. Grids for pressure  $u$  (left picture) and fluxes  $W_1, W_2$  (right picture)

Following notations for the spaces of grid functions defined on  $\omega_0$ ,  $\omega_1$ , and  $\omega_2$  are introduced:

$$H_y = \{y(x), x \in \bar{\omega}_0\}, \quad H_y^0 = \{y(x), x \in \bar{\omega}_0, y(x) = 0 \text{ for } x \in \gamma_0\},$$

$$H_1 = \{w_1(x), x \in \bar{\omega}_1\}, \quad H_2 = \{w_2(x), x \in \bar{\omega}_2\},$$

$$\mathbf{H}_{\mathbf{w}} = H_1 \times H_2, \quad \mathbf{w} = (w_1, w_2).$$

The inner products are defined in the usual way

$$(y, \tilde{y})_{\omega_0} = \sum_{i=1}^{N_1} \sum_{j=1}^{N_2} y_{i,j}(x) \tilde{y}_{i,j}(x) h_1 h_2, \quad \|y\|_{\omega_0} = \sqrt{(y, y)_{\omega_0}}.$$

$$(\mathbf{w}, \tilde{\mathbf{w}})_{\bar{\omega}_1 \times \bar{\omega}_2} = (w_1, \tilde{w}_1)_{\bar{\omega}_1} + (w_2, \tilde{w}_2)_{\bar{\omega}_2} = \sum_{i=0}^{N_1} \sum_{j=1}^{N_2-1} w_1(x) \tilde{w}_1(x) h_i h_j + \sum_{i=1}^{N_1-1} \sum_{j=0}^{N_2} w_2(x) \tilde{w}_2(x) h_i h_j,$$

where  $h_i = h_1$ ,  $i = \overline{1, N_1 - 1}$ ,  $h_0 = h_{N_1} = h_1/2$ ,  $h_j = h_2$ ,  $j = \overline{1, N_2 - 1}$ ,  $h_0 = h_{N_2} = h_2/2$ .

Let us now define discrete operators associated with divergence and gradient operators. Discrete divergence operator  $D : \mathbf{H}_{\mathbf{w}} \rightarrow H_y$  is defined in the following way

$$(D\mathbf{w})_{i,j} = \frac{(w_1)_{i+1,j} - (w_1)_{i,j}}{h_1} + \frac{(w_2)_{i,j+1} - (w_2)_{i,j}}{h_2}, \quad 1 < i \leq N_1, \quad 1 < j \leq N_2.$$



Discrete gradient operator  $G : H_y \rightarrow \mathbf{H}_w$  is defined as  $Gy = (G^1y, G^2y)$ , where

$$G^1y = \begin{cases} \frac{y_{i+1,j} - y_{i,j}}{h_1}, & 1 \leq i \leq N_1 - 1, \\ \frac{y_1 - y_0}{h_1/2}, & i = 0, \\ \frac{y_{N_1+1} - y_{N_1}}{h_1/2}, & i = N_1. \end{cases}$$

$$G^2y = \begin{cases} \frac{y_{i,j+1} - y_{i,j}}{h_2}, & 1 \leq j \leq N_2 - 1, \\ \frac{y_1 - y_0}{h_2/2}, & j = 0, \\ \frac{y_{N_2+1} - y_{N_2}}{h_2/2}, & j = N_2. \end{cases}$$

Note that the above introduced operators satisfy  $D = -G^*$  in the respective scalar products:

$$(Dw, \tilde{y}) = (DGy, \tilde{y})_{\omega_0} = -(Gy, G\tilde{y})_{\omega_1 \times \omega_2}.$$

We will need some auxiliary inequalities, norms, etc., which we list here for completeness.

**Lemma 3.1.** *For any grid function  $y \in H_y^0$  the following inequality (Friedrichs' inequality) holds*

$$\|y\|_{\tilde{\omega}_0} \leq c_1 \|Gy\|_{\tilde{\omega}_1 \times \tilde{\omega}_2}.$$

Discrete  $W_2^1$  norm is defined as

$$\|y\|_{W_2^1} = (y, y)_{\tilde{\omega}_0} + (Gy, Gy)_{\tilde{\omega}_1 \times \tilde{\omega}_2}.$$

From Friedrichs' inequality we get

$$\|y\|_{W_2^1} \leq c_2 \|Gy\|_{\tilde{\omega}_1 \times \tilde{\omega}_2};$$

Let us denote by  $z$  the error,  $z = y - u$ . We will need to write in divergence form the equation for the error. Suppose there exists  $\bar{K}$  such that

$$c_1(w, \tilde{w})_{\omega_1 \times \omega_2} \leq (\bar{K}w, \tilde{w})_{\omega_1 \times \omega_2} \leq c_2(w, \tilde{w})_{\omega_1 \times \omega_2},$$

and such that we can write our discrete equation for the error as

$$-D\bar{K}Gz = -D\psi.$$

Note: Later we will show that for MPFA there exist such  $\bar{K}$  (although we can not write it in explicit form). So, from the inequalities above we have:

$$-(D\bar{K}Gz, z)_{\omega_u} = -(D\psi, z)_{\omega_u} \Rightarrow$$

$$c_1 \|Gz\|_{\omega_1 \times \omega_2}^2 \leq (\bar{K}Gz, Gz)_{\omega_1 \times \omega_2} = (\psi, Gz)_{\omega_1 \times \omega_2} \leq \|\psi\|_{\omega_1 \times \omega_2} \|Gz\|_{\omega_1 \times \omega_2}.$$

Next, we can use the approximation property of MPFA, shown above, to get

$$c_1 \|Gz\|_{\omega_1 \times \omega_2} \leq \|\psi\|_{\omega_1 \times \omega_2} = O(h).$$

This is the end of the proof, because from Friedrichs inequality applied to  $z$  we have the desired result.

It still remains to show that there exist  $\bar{K}$  with the desired properties. From equation  $\tilde{A}\tilde{v} = \tilde{w}$  with  $\tilde{v} = (a^1, b^1, a^3, b^3)^T$ , where  $\tilde{A}$  is given by (9), we see that  $\tilde{\omega}_1, \tilde{\omega}_2, \tilde{\omega}_3, \tilde{\omega}_4$  contain different components of the gradient  $Gy$ , and ONLY components of  $Gy$ . If we denote  $\tilde{B} = \tilde{A}^{-1}$ , we will have

$$a_1 = b_{11}\tilde{w}_1 + b_{12}\tilde{w}_2 + b_{13}\tilde{w}_3 + b_{14}\tilde{w}_4,$$

i.e.,  $a_1$  will also contain only components of  $Gy$ , multiplied by some coefficients independent on the solution. Similar conclusion can be drawn for  $b_1, a_2, b_2$ , and

other similar expressions which we use to write our finite difference scheme 11. Note, that 11 is already written in divergence form, thus we are able to write MPFA as  $D\bar{K}Gy$ , with some  $\bar{K}$  which can not be explicitly written. Next, we have to clarify what are the properties of  $\bar{K}$ . From [1] we know that on rectangular grid MPFA can be written as  $My = f$  with  $M = M^* > 0$ . Thus we have

$$My = f, \quad -D\bar{K}Gy = f, \Rightarrow M = D\bar{K}G.$$

It follows from here that

$$(My, y) = -(D\bar{K}Gy, y) = -(\bar{K}Gy, Gy).$$

Now we can conclude: i) Because  $M$  is symmetric, it follows  $\bar{K}$  is also symmetric. ii) Because  $(My, y) \geq 0$ ,  $\Rightarrow (\bar{K}Gy, Gy) \geq 0$ . Moreover,  $(My, y) > 0$  for  $y \neq 0$ . It remains to answer the question is  $(\bar{K}Gy, Gy) > 0$  for  $Gy \neq 0$ ? In our case,  $y = const \Rightarrow$  is possible only for  $y \equiv 0$ , and with this we have shown that there exist  $\bar{K}$  with the desired properties.

#### 4. NUMERICAL RESULTS

In this section, we validate the developed difference scheme by performing several numerical experiments for variable continuous and discontinuous full tensor coefficients.

**4.1. Validation of the discretization for continuous coefficients.** Consider the example from [3]. In this experiment, the permeability tensor is full symmetrical with variable continuous components

$$(27) \quad K = \begin{pmatrix} (x+2)^2 + y^2 & \sin(xy) \\ \sin(xy) & 1 \end{pmatrix}$$

The exact solution is given by the following formula

$$u = x^3y + y^4 + \sin(x) \cos(y),$$

with  $f$  and  $g^D$  defined accordingly by (1), (2).

Convergence rate for the pressure is established by running cases for seven levels of grid refinement, starting with  $5 \times 5$  grid cells on level one and refining the grid steps by a factor of two for each successive level. Further, we will use the following notations

$$\|\delta_u\|_C = \frac{\|u - u_h\|_C}{\|u\|_C}, \quad \|\delta_u\|_{L_2} = \frac{\|u - u_h\|_{L_2}}{\|u\|_{L_2}},$$

where  $u$  is the solution of the differential problem (1), (2), and  $u_h$  is the solution of the difference scheme (11).

From Tab. 1 we see that if we refine the grid step by the factor of two, then the error reduces by the factor of four. Hence, the approximate solution  $u_h$  converges to the exact solution  $u$  with the second order both in the maximum and  $L_2$  norms.

In the same way we calculate the convergence for the fluxes  $W_1$  and  $W_2$ . In Tab. 2 the following notations are used

$$\begin{aligned} \|\delta_1\|_C &= \frac{\|W_1 - W_{1,h}\|_C}{\|W_1\|_C}, & \|\delta_1\|_{L_2} &= \frac{\|W_1 - W_{1,h}\|_{L_2}}{\|W_1\|_{L_2}}, \\ \|\delta_2\|_C &= \frac{\|W_2 - W_{2,h}\|_C}{\|W_2\|_C}, & \|\delta_2\|_{L_2} &= \frac{\|W_2 - W_{2,h}\|_{L_2}}{\|W_2\|_{L_2}}. \end{aligned}$$

Grid	$\ \delta_u\ _C$	$\ \delta_u\ _{L_2}$
$5 \times 5$	$1.97 \times 10^{-2}$	$1.73 \times 10^{-2}$
$10 \times 10$	$6.02 \times 10^{-3}$	$4.87 \times 10^{-3}$
$20 \times 20$	$1.56 \times 10^{-3}$	$1.23 \times 10^{-3}$
$40 \times 40$	$3.84 \times 10^{-4}$	$3.05 \times 10^{-4}$
$80 \times 80$	$9.49 \times 10^{-5}$	$7.56 \times 10^{-5}$
$160 \times 160$	$2.36 \times 10^{-5}$	$1.86 \times 10^{-5}$
$320 \times 320$	$5.90 \times 10^{-6}$	$4.64 \times 10^{-6}$

TABLE 1. Convergence of the pressure for problem (1), (2) with permeability tensor (27)

Grid	$\ \delta_1\ _C$	$\ \delta_1\ _{L_2}$	$\ \delta_2\ _C$	$\ \delta_2\ _{L_2}$
$5 \times 5$	$7.78 \times 10^{-3}$	$7.76 \times 10^{-3}$	$7.47 \times 10^{-3}$	$6.83 \times 10^{-3}$
$10 \times 10$	$4.69 \times 10^{-3}$	$2.98 \times 10^{-3}$	$4.76 \times 10^{-3}$	$4.18 \times 10^{-3}$
$20 \times 20$	$2.21 \times 10^{-3}$	$9.93 \times 10^{-4}$	$2.20 \times 10^{-3}$	$1.73 \times 10^{-3}$
$40 \times 40$	$1.00 \times 10^{-3}$	$2.88 \times 10^{-4}$	$9.23 \times 10^{-4}$	$5.40 \times 10^{-4}$
$80 \times 80$	$4.72 \times 10^{-4}$	$8.08 \times 10^{-5}$	$4.29 \times 10^{-4}$	$1.56 \times 10^{-4}$
$160 \times 160$	$2.27 \times 10^{-4}$	$2.23 \times 10^{-5}$	$2.09 \times 10^{-4}$	$4.39 \times 10^{-5}$
$320 \times 320$	$1.11 \times 10^{-4}$	$6.09 \times 10^{-6}$	$1.03 \times 10^{-4}$	$1.21 \times 10^{-5}$

TABLE 2. Convergence of the fluxes for problem (1), (2) with permeability tensor (27)

From Tab. 2 we can see that for the fluxes we have the first order convergence in the maximum norm and the second order convergence in the  $L_2$  norm.

**4.2. Validation of the discretization for discontinuous coefficients.** Consider jump discontinuity on the interfaces  $x_\xi = 0.4$  and  $y_\eta = 0.4$ . The permeability tensor is given by

$$(28) \quad K_1 = K_4 = \begin{pmatrix} 1 & 1/2 \\ 1/2 & 1 \end{pmatrix}, \quad K_2 = K_3 = \begin{pmatrix} \lambda & \lambda/2 \\ \lambda/2 & \lambda \end{pmatrix},$$

where  $\lambda$  is called the contrast of discontinuity. We choose the exact solution which satisfies the conditions of the pressure and fluxes continuity across the interfaces

$$u = (x - x_\xi)^2 (y - y_\eta)^2 \sin(\pi(x + y)).$$

The right hand side and the boundary conditions can be found from the initial differential problem (1), (2) for known exact solution. The experiments are performed for  $\lambda = 10$  and  $\lambda = 1000$ .

From Tab. 3 it is easy to see that the approximate solution  $u_h$  converges to the exact solution  $u$  with the second order both in the maximum and  $L_2$  norms. For the fluxes we have first order convergence in the  $C$  norm and the second order convergence in the  $L_2$  norm (see Tab. 4).

Note, that for this artificial smooth solution the accuracy does not depend on jump discontinuity.




FIGURE 3. The surface of discontinuity

Grid	$\ \delta_u\ _C$		$\ \delta_u\ _{L_2}$	
	$\lambda = 10$	$\lambda = 1000$	$\lambda = 10$	$\lambda = 10000$
$5 \times 5$	$1.56 \times 10^{-1}$	$1.56 \times 10^{-1}$	$1.61 \times 10^{-1}$	$1.63 \times 10^{-1}$
$10 \times 10$	$5.77 \times 10^{-2}$	$5.78 \times 10^{-2}$	$4.58 \times 10^{-2}$	$4.64 \times 10^{-2}$
$20 \times 20$	$1.72 \times 10^{-2}$	$1.73 \times 10^{-2}$	$1.18 \times 10^{-2}$	$1.20 \times 10^{-2}$
$40 \times 40$	$4.91 \times 10^{-3}$	$4.91 \times 10^{-3}$	$3.02 \times 10^{-3}$	$3.06 \times 10^{-3}$
$80 \times 80$	$1.30 \times 10^{-3}$	$1.30 \times 10^{-3}$	$7.67 \times 10^{-4}$	$7.77 \times 10^{-4}$
$160 \times 160$	$3.37 \times 10^{-4}$	$3.37 \times 10^{-4}$	$1.93 \times 10^{-4}$	$1.96 \times 10^{-4}$
$320 \times 320$	$8.66 \times 10^{-5}$	$8.66 \times 10^{-5}$	$4.86 \times 10^{-5}$	$4.92 \times 10^{-5}$

TABLE 3. Convergence of the pressure for problem (1), (2) with permeability tensor (28)

Grid	$\ \delta_1\ _C = \ \delta_2\ _C$		$\ \delta_1\ _{L_2} = \ \delta_2\ _{L_2}$	
	$\lambda = 10$	$\lambda = 1000$	$\lambda = 10$	$\lambda = 10000$
$5 \times 5$	$1.63 \times 10^{-1}$	$1.72 \times 10^{-1}$	$1.92 \times 10^{-1}$	$2.29 \times 10^{-1}$
$10 \times 10$	$1.51 \times 10^{-2}$	$1.60 \times 10^{-2}$	$2.92 \times 10^{-2}$	$3.06 \times 10^{-2}$
$20 \times 20$	$1.70 \times 10^{-2}$	$1.70 \times 10^{-2}$	$1.05 \times 10^{-2}$	$1.08 \times 10^{-2}$
$40 \times 40$	$1.04 \times 10^{-2}$	$1.04 \times 10^{-2}$	$4.32 \times 10^{-3}$	$4.39 \times 10^{-3}$
$80 \times 80$	$5.59 \times 10^{-3}$	$5.59 \times 10^{-3}$	$1.52 \times 10^{-3}$	$1.54 \times 10^{-3}$
$160 \times 160$	$2.87 \times 10^{-3}$	$2.87 \times 10^{-3}$	$4.89 \times 10^{-4}$	$4.95 \times 10^{-4}$
$320 \times 320$	$1.45 \times 10^{-3}$	$1.45 \times 10^{-3}$	$1.47 \times 10^{-4}$	$1.49 \times 10^{-4}$

TABLE 4. Convergence of the fluxes for problem (1), (2) with permeability tensor (28)

## 5. CONCLUDING REMARKS

The purpose of the paper was to study the approximation properties of the multipoint flux approximation discretization for two-dimensional second order elliptic equations in the case of discontinuous tensor coefficients. The discretization is based on the finite volume method and multipoint flux approximation approach. Specific derivation of MPFA is written and it is shown that the components of the discrete flux approximate the components of the continuous flux with the first order in the midpoints of the edges. The error of the finite difference scheme is written in divergence form, so this is used to show first order convergence for the fluxes in

discrete  $W_2^1$  norm. The analysis is also valid for the problems with continuous tensor coefficients. Results from numerical experiments are presented and discussed.

#### ACKNOWLEDGMENTS

This work has been partially supported by EC under the project INTAS-30-50-4395, by Deutscher Akademischer Austausch Dienst (I. Rybak, scholarship grant A/05/57218) and by Belarusian Republican Foundation for Fundamental Research (project F07MS-054).

#### REFERENCES

- [1] Aavatsmark I. An introduction to multipoint flux approximations for quadrilateral grids, *Comput. Geosci.*, 6(2002), pp. 405–432.
- [2] Aavatsmark I., Eigestad G.T., Klausen R.A., Wheeler M.F., and Yotov I. Convergence of a symmetric MPFA method on quadrilateral grids, *Comput. Geosci.*, submitted.
- [3] Arbogast T., Wheeler M.F., Yotov I., Mixed finite elements for elliptic problems with tensor coefficients as cell-centered finite differences, *SIAM J. Numer. Anal.*, 34(1997), No.2, pp. 828–852.
- [4] Eigestad G.T., Klausen R.A. On the convergence of the multi-point flux approximation O-method: numerical experiments for discontinuous permeability, *Numer. Methods Partial Diff. Eqns.*, 21(2005), No.6, pp. 1079–1098.
- [5] Klausen R.A., Winther R. Convergence of multipoint flux approximations on quadrilateral grids, *Numer. Methods Partial Diff. Eqns.*, 22(2006), No.6, pp. 1438–1454.
- [6] Petzoldt, M., A posteriori error estimators for elliptic equations with discontinuous diffusion coefficients, *WIAS Preprint No. 533*, (1999) <http://www.wias-berlin.de/main/publications/wias-publ/>
- [7] Samarskii A. A., *The Theory of Difference Schemes*, Marcel Dekker, Inc., New York–Basel, 2001.
- [8] Ware A.F., Parrott A.K., Rogers C., A finite volume discretisation for porous media flows governed by non-diagonal permeability tensors, *Proceedings of CFD95, Third Annual Conference of the CFD Society of Canada, Banff, Alberta, Canada, 25–27 June, 1995*.
- [9] Wheeler M., Yotov I. A multipoint flux mixed finite element method, *SIAM J. Numer. Anal.*, 4 (2006), No.5, pp. 2082–2106.
- [10] Edwards M.G., Rogers C.F. Finite volume discretization with imposed flux continuity for the general tensor pressure equation, *Comput. Geosci.*, 2(1998), pp. 259–290.
- [11] Samarskii A. A., Andreev V.B. *Difference Methods for Elliptic Equations*, (In Russian), Nauka, Moscow, 1976.
- [12] Samarskii A. A., Lazarov R.D., Makarov V.L. *Difference Schemes for Differential Equations having Generalized Solutions*, (In Russian), Vysshaya Shkola Publishers, Moscow, 1987.

FRAUNHOFER INSTITUT FUER TECHNO- UND WIRTSCHAFTSMATHEMATIK, FRAUNHOFER-PLATZ 1, 67663 KAISERSLAUTERN, GERMANY, AND INSTITUTE OF MATHEMATICS, BULGARIAN ACADEMY OF SCIENCE, ACAD. G.BONCHEV STR., BL.8, BG-1113, SOFIA, BULGARIA, ILIEV@ITWM.FHG.DE

INSTITUTE OF MATHEMATICS, NATIONAL ACADEMY OF SCIENCES OF BELARUS, SURGANOV STR. 11, 220072 MINSK, BELARUS, RYBAK@IM.BAS-NET.BY

# Published reports of the Fraunhofer ITWM

The PDF-files of the following reports are available under:

[www.itwm.fraunhofer.de/de/zentral\\_\\_berichte/berichte](http://www.itwm.fraunhofer.de/de/zentral__berichte/berichte)

1. D. Hietel, K. Steiner, J. Struckmeier  
**A Finite - Volume Particle Method for Compressible Flows**  
(19 pages, 1998)
2. M. Feldmann, S. Seibold  
**Damage Diagnosis of Rotors: Application of Hilbert Transform and Multi-Hypothesis Testing**  
*Keywords: Hilbert transform, damage diagnosis, Kalman filtering, non-linear dynamics*  
(23 pages, 1998)
3. Y. Ben-Haim, S. Seibold  
**Robust Reliability of Diagnostic Multi-Hypothesis Algorithms: Application to Rotating Machinery**  
*Keywords: Robust reliability, convex models, Kalman filtering, multi-hypothesis diagnosis, rotating machinery, crack diagnosis*  
(24 pages, 1998)
4. F.-Th. Lentjes, N. Siedow  
**Three-dimensional Radiative Heat Transfer in Glass Cooling Processes**  
(23 pages, 1998)
5. A. Klar, R. Wegener  
**A hierarchy of models for multilane vehicular traffic**  
**Part I: Modeling**  
(23 pages, 1998)  
  
**Part II: Numerical and stochastic investigations**  
(17 pages, 1998)
6. A. Klar, N. Siedow  
**Boundary Layers and Domain Decomposition for Radiative Heat Transfer and Diffusion Equations: Applications to Glass Manufacturing Processes**  
(24 pages, 1998)
7. I. Choquet  
**Heterogeneous catalysis modelling and numerical simulation in rarified gas flows**  
**Part I: Coverage locally at equilibrium**  
(24 pages, 1998)
8. J. Ohser, B. Steinbach, C. Lang  
**Efficient Texture Analysis of Binary Images**  
(17 pages, 1998)
9. J. Orlik  
**Homogenization for viscoelasticity of the integral type with aging and shrinkage**  
(20 pages, 1998)
10. J. Mohring  
**Helmholtz Resonators with Large Aperture**  
(21 pages, 1998)
11. H. W. Hamacher, A. Schöbel  
**On Center Cycles in Grid Graphs**  
(15 pages, 1998)
12. H. W. Hamacher, K.-H. Küfer  
**Inverse radiation therapy planning - a multiple objective optimisation approach**  
(14 pages, 1999)
13. C. Lang, J. Ohser, R. Hilfer  
**On the Analysis of Spatial Binary Images**  
(20 pages, 1999)
14. M. Junk  
**On the Construction of Discrete Equilibrium Distributions for Kinetic Schemes**  
(24 pages, 1999)
15. M. Junk, S. V. Raghurame Rao  
**A new discrete velocity method for Navier-Stokes equations**  
(20 pages, 1999)
16. H. Neunzert  
**Mathematics as a Key to Key Technologies**  
(39 pages (4 PDF-Files), 1999)
17. J. Ohser, K. Sandau  
**Considerations about the Estimation of the Size Distribution in Wicksell's Corpuscle Problem**  
(18 pages, 1999)
18. E. Carrizosa, H. W. Hamacher, R. Klein, S. Nickel  
**Solving nonconvex planar location problems by finite dominating sets**  
*Keywords: Continuous Location, Polyhedral Gauges, Finite Dominating Sets, Approximation, Sandwich Algorithm, Greedy Algorithm*  
(19 pages, 2000)
19. A. Becker  
**A Review on Image Distortion Measures**  
*Keywords: Distortion measure, human visual system*  
(26 pages, 2000)
20. H. W. Hamacher, M. Labbé, S. Nickel, T. Sonneborn  
**Polyhedral Properties of the Uncapacitated Multiple Allocation Hub Location Problem**  
*Keywords: integer programming, hub location, facility location, valid inequalities, facets, branch and cut*  
(21 pages, 2000)
21. H. W. Hamacher, A. Schöbel  
**Design of Zone Tariff Systems in Public Transportation**  
(30 pages, 2001)
22. D. Hietel, M. Junk, R. Keck, D. Teleaga  
**The Finite-Volume-Particle Method for Conservation Laws**  
(16 pages, 2001)
23. T. Bender, H. Hennes, J. Kalcsics, M. T. Melo, S. Nickel  
**Location Software and Interface with GIS and Supply Chain Management**  
*Keywords: facility location, software development, geographical information systems, supply chain management*  
(48 pages, 2001)
24. H. W. Hamacher, S. A. Tjandra  
**Mathematical Modelling of Evacuation Problems: A State of Art**  
(44 pages, 2001)
25. J. Kuhnert, S. Tiwari  
**Grid free method for solving the Poisson equation**  
*Keywords: Poisson equation, Least squares method, Grid free method*  
(19 pages, 2001)
26. T. Götz, H. Rave, D. Reinel-Bitzer, K. Steiner, H. Tiemeier  
**Simulation of the fiber spinning process**  
*Keywords: Melt spinning, fiber model, Lattice Boltzmann, CFD*  
(19 pages, 2001)
27. A. Zemitis  
**On interaction of a liquid film with an obstacle**  
*Keywords: impinging jets, liquid film, models, numerical solution, shape*  
(22 pages, 2001)
28. I. Ginzburg, K. Steiner  
**Free surface lattice-Boltzmann method to model the filling of expanding cavities by Bingham Fluids**  
*Keywords: Generalized LBE, free-surface phenomena, interface boundary conditions, filling processes, Bingham viscoplastic model, regularized models*  
(22 pages, 2001)
29. H. Neunzert  
**»Denn nichts ist für den Menschen als Menschen etwas wert, was er nicht mit Leidenschaft tun kann«**  
**Vortrag anlässlich der Verleihung des Akademiepreises des Landes Rheinland-Pfalz am 21.11.2001**  
*Keywords: Lehre, Forschung, angewandte Mathematik, Mehrskalalanalyse, Strömungsmechanik*  
(18 pages, 2001)
30. J. Kuhnert, S. Tiwari  
**Finite pointset method based on the projection method for simulations of the incompressible Navier-Stokes equations**  
*Keywords: Incompressible Navier-Stokes equations, Meshfree method, Projection method, Particle scheme, Least squares approximation*  
*AMS subject classification: 76D05, 76M28*  
(25 pages, 2001)
31. R. Korn, M. Krekel  
**Optimal Portfolios with Fixed Consumption or Income Streams**  
*Keywords: Portfolio optimisation, stochastic control, HJB equation, discretisation of control problems.*  
(23 pages, 2002)
32. M. Krekel  
**Optimal portfolios with a loan dependent credit spread**  
*Keywords: Portfolio optimisation, stochastic control, HJB equation, credit spread, log utility, power utility, non-linear wealth dynamics*  
(25 pages, 2002)
33. J. Ohser, W. Nagel, K. Schladitz  
**The Euler number of discretized sets – on the choice of adjacency in homogeneous lattices**  
*Keywords: image analysis, Euler number, neighborhood relationships, cuboidal lattice*  
(32 pages, 2002)



34. I. Ginzburg, K. Steiner  
**Lattice Boltzmann Model for Free-Surface flow and Its Application to Filling Process in Casting**  
Keywords: Lattice Boltzmann models; free-surface phenomena; interface boundary conditions; filling processes; injection molding; volume of fluid method; interface boundary conditions; advection-schemes; up-wind-schemes (54 pages, 2002)
35. M. Günther, A. Klar, T. Materne, R. Wegener  
**Multivalued fundamental diagrams and stop and go waves for continuum traffic equations**  
Keywords: traffic flow, macroscopic equations, kinetic derivation, multivalued fundamental diagram, stop and go waves, phase transitions (25 pages, 2002)
36. S. Feldmann, P. Lang, D. Prätzel-Wolters  
**Parameter influence on the zeros of network determinants**  
Keywords: Networks, Equicofactor matrix polynomials, Realization theory, Matrix perturbation theory (30 pages, 2002)
37. K. Koch, J. Ohser, K. Schladitz  
**Spectral theory for random closed sets and estimating the covariance via frequency space**  
Keywords: Random set, Bartlett spectrum, fast Fourier transform, power spectrum (28 pages, 2002)
38. D. d'Humières, I. Ginzburg  
**Multi-reflection boundary conditions for lattice Boltzmann models**  
Keywords: lattice Boltzmann equation, boundary conditions, bounce-back rule, Navier-Stokes equation (72 pages, 2002)
39. R. Korn  
**Elementare Finanzmathematik**  
Keywords: Finanzmathematik, Aktien, Optionen, Portfolio-Optimierung, Börse, Lehrerweiterbildung, Mathematikunterricht (98 pages, 2002)
40. J. Kallrath, M. C. Müller, S. Nickel  
**Batch Presorting Problems: Models and Complexity Results**  
Keywords: Complexity theory, Integer programming, Assignment, Logistics (19 pages, 2002)
41. J. Linn  
**On the frame-invariant description of the phase space of the Folgar-Tucker equation**  
Key words: fiber orientation, Folgar-Tucker equation, injection molding (5 pages, 2003)
42. T. Hanne, S. Nickel  
**A Multi-Objective Evolutionary Algorithm for Scheduling and Inspection Planning in Software Development Projects**  
Key words: multiple objective programming, project management and scheduling, software development, evolutionary algorithms, efficient set (29 pages, 2003)
43. T. Bortfeld, K.-H. Küfer, M. Monz, A. Scherrer, C. Thieke, H. Trinkaus  
**Intensity-Modulated Radiotherapy - A Large Scale Multi-Criteria Programming Problem**  
Keywords: multiple criteria optimization, representative systems of Pareto solutions, adaptive triangulation, clustering and disaggregation techniques, visualization of Pareto solutions, medical physics, external beam radiotherapy planning, intensity modulated radiotherapy (31 pages, 2003)
44. T. Halfmann, T. Wichmann  
**Overview of Symbolic Methods in Industrial Analog Circuit Design**  
Keywords: CAD, automated analog circuit design, symbolic analysis, computer algebra, behavioral modeling, system simulation, circuit sizing, macro modeling, differential-algebraic equations, index (17 pages, 2003)
45. S. E. Mikhailov, J. Orlik  
**Asymptotic Homogenisation in Strength and Fatigue Durability Analysis of Composites**  
Keywords: multiscale structures, asymptotic homogenization, strength, fatigue, singularity, non-local conditions (14 pages, 2003)
46. P. Domínguez-Marín, P. Hansen, N. Mladenović, S. Nickel  
**Heuristic Procedures for Solving the Discrete Ordered Median Problem**  
Keywords: genetic algorithms, variable neighborhood search, discrete facility location (31 pages, 2003)
47. N. Boland, P. Domínguez-Marín, S. Nickel, J. Puerto  
**Exact Procedures for Solving the Discrete Ordered Median Problem**  
Keywords: discrete location, Integer programming (41 pages, 2003)
48. S. Feldmann, P. Lang  
**Padé-like reduction of stable discrete linear systems preserving their stability**  
Keywords: Discrete linear systems, model reduction, stability, Hankel matrix, Stein equation (16 pages, 2003)
49. J. Kallrath, S. Nickel  
**A Polynomial Case of the Batch Presorting Problem**  
Keywords: batch presorting problem, online optimization, competitive analysis, polynomial algorithms, logistics (17 pages, 2003)
50. T. Hanne, H. L. Trinkaus  
**knowCube for MCDM – Visual and Interactive Support for Multicriteria Decision Making**  
Key words: Multicriteria decision making, knowledge management, decision support systems, visual interfaces, interactive navigation, real-life applications. (26 pages, 2003)
51. O. Iliev, V. Laptev  
**On Numerical Simulation of Flow Through Oil Filters**  
Keywords: oil filters, coupled flow in plain and porous media, Navier-Stokes, Brinkman, numerical simulation (8 pages, 2003)
52. W. Dörfler, O. Iliev, D. Stoyanov, D. Vassileva  
**On a Multigrid Adaptive Refinement Solver for Saturated Non-Newtonian Flow in Porous Media**  
Keywords: Nonlinear multigrid, adaptive refinement, Heston model, stochastic volatility, cliquet options (17 pages, 2003)
53. S. Kruse  
**On the Pricing of Forward Starting Options under Stochastic Volatility**  
Keywords: Option pricing, forward starting options, Heston model, stochastic volatility, cliquet options (11 pages, 2003)
54. O. Iliev, D. Stoyanov  
**Multigrid – adaptive local refinement solver for incompressible flows**  
Keywords: Navier-Stokes equations, incompressible flow, projection-type splitting, SIMPLE, multigrid methods, adaptive local refinement, lid-driven flow in a cavity (37 pages, 2003)
55. V. Starikovicius  
**The multiphase flow and heat transfer in porous media**  
Keywords: Two-phase flow in porous media, various formulations, global pressure, multiphase mixture model, numerical simulation (30 pages, 2003)
56. P. Lang, A. Sarishvili, A. Wirsén  
**Blocked neural networks for knowledge extraction in the software development process**  
Keywords: Blocked Neural Networks, Nonlinear Regression, Knowledge Extraction, Code Inspection (21 pages, 2003)
57. H. Knaf, P. Lang, S. Zeiser  
**Diagnosis aiding in Regulation Thermography using Fuzzy Logic**  
Keywords: fuzzy logic, knowledge representation, expert system (22 pages, 2003)
58. M. T. Melo, S. Nickel, F. Saldanha da Gama  
**Largescale models for dynamic multi-commodity capacitated facility location**  
Keywords: supply chain management, strategic planning, dynamic location, modeling (40 pages, 2003)
59. J. Orlik  
**Homogenization for contact problems with periodically rough surfaces**  
Keywords: asymptotic homogenization, contact problems (28 pages, 2004)
60. A. Scherrer, K.-H. Küfer, M. Monz, F. Alonso, T. Bortfeld  
**IMRT planning on adaptive volume structures – a significant advance of computational complexity**  
Keywords: Intensity-modulated radiation therapy (IMRT), inverse treatment planning, adaptive volume structures, hierarchical clustering, local refinement, adaptive clustering, convex programming, mesh generation, multi-grid methods (24 pages, 2004)

61. D. Kehrwald  
**Parallel lattice Boltzmann simulation of complex flows**  
Keywords: Lattice Boltzmann methods, parallel computing, microstructure simulation, virtual material design, pseudo-plastic fluids, liquid composite moulding (12 pages, 2004)
62. O. Iliev, J. Linn, M. Moog, D. Niedziela, V. Starikovicus  
**On the Performance of Certain Iterative Solvers for Coupled Systems Arising in Discretization of Non-Newtonian Flow Equations**  
Keywords: Performance of iterative solvers, Preconditioners, Non-Newtonian flow (17 pages, 2004)
63. R. Ciegis, O. Iliev, S. Rief, K. Steiner  
**On Modelling and Simulation of Different Regimes for Liquid Polymer Moulding**  
Keywords: Liquid Polymer Moulding, Modelling, Simulation, Infiltration, Front Propagation, non-Newtonian flow in porous media (43 pages, 2004)
64. T. Hanne, H. Neu  
**Simulating Human Resources in Software Development Processes**  
Keywords: Human resource modeling, software process, productivity, human factors, learning curve (14 pages, 2004)
65. O. Iliev, A. Mikelic, P. Popov  
**Fluid structure interaction problems in deformable porous media: Toward permeability of deformable porous media**  
Keywords: fluid-structure interaction, deformable porous media, upscaling, linear elasticity, stokes, finite elements (28 pages, 2004)
66. F. Gaspar, O. Iliev, F. Lisbona, A. Naumovich, P. Vabishchevich  
**On numerical solution of 1-D poroelasticity equations in a multilayered domain**  
Keywords: poroelasticity, multilayered material, finite volume discretization, MAC type grid (41 pages, 2004)
67. J. Ohser, K. Schladitz, K. Koch, M. Nöthe  
**Diffraction by image processing and its application in materials science**  
Keywords: porous microstructure, image analysis, random set, fast Fourier transform, power spectrum, Bartlett spectrum (13 pages, 2004)
68. H. Neunzert  
**Mathematics as a Technology: Challenges for the next 10 Years**  
Keywords: applied mathematics, technology, modelling, simulation, visualization, optimization, glass processing, spinning processes, fiber-fluid interaction, turbulence effects, topological optimization, multicriteria optimization, Uncertainty and Risk, financial mathematics, Malliavin calculus, Monte-Carlo methods, virtual material design, filtration, bio-informatics, system biology (29 pages, 2004)
69. R. Ewing, O. Iliev, R. Lazarov, A. Naumovich  
**On convergence of certain finite difference discretizations for 1D poroelasticity interface problems**  
Keywords: poroelasticity, multilayered material, finite volume discretizations, MAC type grid, error estimates (26 pages, 2004)
70. W. Dörfler, O. Iliev, D. Stoyanov, D. Vassileva  
**On Efficient Simulation of Non-Newtonian Flow in Saturated Porous Media with a Multigrid Adaptive Refinement Solver**  
Keywords: Nonlinear multigrid, adaptive refinement, non-Newtonian in porous media (25 pages, 2004)
71. J. Kalcsics, S. Nickel, M. Schröder  
**Towards a Unified Territory Design Approach – Applications, Algorithms and GIS Integration**  
Keywords: territory design, political districting, sales territory alignment, optimization algorithms, Geographical Information Systems (40 pages, 2005)
72. K. Schladitz, S. Peters, D. Reinel-Bitzer, A. Wiegmann, J. Ohser  
**Design of acoustic trim based on geometric modeling and flow simulation for non-woven**  
Keywords: random system of fibers, Poisson line process, flow resistivity, acoustic absorption, Lattice-Boltzmann method, non-woven (21 pages, 2005)
73. V. Rutka, A. Wiegmann  
**Explicit Jump Immersed Interface Method for virtual material design of the effective elastic moduli of composite materials**  
Keywords: virtual material design, explicit jump immersed interface method, effective elastic moduli, composite materials (22 pages, 2005)
74. T. Hanne  
**Eine Übersicht zum Scheduling von Baustellen**  
Keywords: Projektplanung, Scheduling, Bauplanung, Bauindustrie (32 pages, 2005)
75. J. Linn  
**The Folgar-Tucker Model as a Differential Algebraic System for Fiber Orientation Calculation**  
Keywords: fiber orientation, Folgar-Tucker model, invariants, algebraic constraints, phase space, trace stability (15 pages, 2005)
76. M. Speckert, K. Dreßler, H. Mauch, A. Lion, G. J. Wierda  
**Simulation eines neuartigen Prüfsystems für Achserproben durch MKS-Modellierung einschließlich Regelung**  
Keywords: virtual test rig, suspension testing, multi-body simulation, modeling hexapod test rig, optimization of test rig configuration (20 pages, 2005)
77. K.-H. Küfer, M. Monz, A. Scherrer, P. Süß, F. Alonso, A. S. A. Sultan, Th. Bortfeld, D. Craft, Chr. Thieke  
**Multicriteria optimization in intensity modulated radiotherapy planning**  
Keywords: multicriteria optimization, extreme solutions, real-time decision making, adaptive approximation schemes, clustering methods, IMRT planning, reverse engineering (51 pages, 2005)
78. S. Amstutz, H. Andrä  
**A new algorithm for topology optimization using a level-set method**  
Keywords: shape optimization, topology optimization, topological sensitivity, level-set (22 pages, 2005)
79. N. Ettrich  
**Generation of surface elevation models for urban drainage simulation**  
Keywords: Flooding, simulation, urban elevation models, laser scanning (22 pages, 2005)
80. H. Andrä, J. Linn, I. Matei, I. Shklyar, K. Steiner, E. Teichmann  
**OPTCAST – Entwicklung adäquater Strukturoptimierungsverfahren für Gießereien Technischer Bericht (KURZFASSUNG)**  
Keywords: Topologieoptimierung, Level-Set-Methode, Gießprozesssimulation, Gießtechnische Restriktionen, CAE-Kette zur Strukturoptimierung (77 pages, 2005)
81. N. Marheineke, R. Wegener  
**Fiber Dynamics in Turbulent Flows Part I: General Modeling Framework**  
Keywords: fiber-fluid interaction; Cosserat rod; turbulence modeling; Kolmogorov's energy spectrum; double-velocity correlations; differentiable Gaussian fields (20 pages, 2005)  
**Part II: Specific Taylor Drag**  
Keywords: flexible fibers;  $k-\epsilon$  turbulence model; fiber-turbulence interaction scales; air drag; random Gaussian aerodynamic force; white noise; stochastic differential equations; ARMA process (18 pages, 2005)
82. C. H. Lampert, O. Wirjadi  
**An Optimal Non-Orthogonal Separation of the Anisotropic Gaussian Convolution Filter**  
Keywords: Anisotropic Gaussian filter, linear filtering, orientation space, nD image processing, separable filters (25 pages, 2005)
83. H. Andrä, D. Stoyanov  
**Error indicators in the parallel finite element solver for linear elasticity DDFEM**  
Keywords: linear elasticity, finite element method, hierarchical shape functions, domain decomposition, parallel implementation, a posteriori error estimates (21 pages, 2006)
84. M. Schröder, I. Solchenbach  
**Optimization of Transfer Quality in Regional Public Transit**  
Keywords: public transit, transfer quality, quadratic assignment problem (16 pages, 2006)
85. A. Naumovich, F. J. Gaspar  
**On a multigrid solver for the three-dimensional Biot poroelasticity system in multilayered domains**  
Keywords: poroelasticity, interface problem, multigrid, operator-dependent prolongation (11 pages, 2006)
86. S. Panda, R. Wegener, N. Marheineke  
**Slender Body Theory for the Dynamics of Curved Viscous Fibers**  
Keywords: curved viscous fibers; fluid dynamics; Navier-Stokes equations; free boundary value problem; asymptotic expansions; slender body theory (14 pages, 2006)
87. E. Ivanov, H. Andrä, A. Kudryavtsev  
**Domain Decomposition Approach for Automatic Parallel Generation of Tetrahedral Grids**  
Key words: Grid Generation, Unstructured Grid, Delaunay Triangulation, Parallel Programming, Domain Decomposition, Load Balancing (18 pages, 2006)



88. S. Tiwari, S. Antonov, D. Hietel, J. Kuhnert, R. Wegener  
**A Meshfree Method for Simulations of Interactions between Fluids and Flexible Structures**  
Key words: Meshfree Method, FPM, Fluid Structure Interaction, Sheet of Paper, Dynamical Coupling (16 pages, 2006)
89. R. Ciegis, O. Iliev, V. Starikovicius, K. Steiner  
**Numerical Algorithms for Solving Problems of Multiphase Flows in Porous Media**  
Keywords: nonlinear algorithms, finite-volume method, software tools, porous media, flows (16 pages, 2006)
90. D. Niedziela, O. Iliev, A. Latz  
**On 3D Numerical Simulations of Viscoelastic Fluids**  
Keywords: non-Newtonian fluids, anisotropic viscosity, integral constitutive equation (18 pages, 2006)
91. A. Winterfeld  
**Application of general semi-infinite Programming to Lapidary Cutting Problems**  
Keywords: large scale optimization, nonlinear programming, general semi-infinite optimization, design centering, clustering (26 pages, 2006)
92. J. Orlik, A. Ostrovska  
**Space-Time Finite Element Approximation and Numerical Solution of Hereditary Linear Viscoelasticity Problems**  
Keywords: hereditary viscoelasticity; kern approximation by interpolation; space-time finite element approximation, stability and a priori estimate (24 pages, 2006)
93. V. Rutka, A. Wiegmann, H. Andrä  
**EJIM for Calculation of effective Elastic Moduli in 3D Linear Elasticity**  
Keywords: Elliptic PDE, linear elasticity, irregular domain, finite differences, fast solvers, effective elastic moduli (24 pages, 2006)
94. A. Wiegmann, A. Zemitis  
**EJ-HEAT: A Fast Explicit Jump Harmonic Averaging Solver for the Effective Heat Conductivity of Composite Materials**  
Keywords: Stationary heat equation, effective thermal conductivity, explicit jump, discontinuous coefficients, virtual material design, microstructure simulation, EJ-HEAT (21 pages, 2006)
95. A. Naumovich  
**On a finite volume discretization of the three-dimensional Biot poroelasticity system in multilayered domains**  
Keywords: Biot poroelasticity system, interface problems, finite volume discretization, finite difference method. (21 pages, 2006)
96. M. Krekel, J. Wenzel  
**A unified approach to Credit Default Swaption and Constant Maturity Credit Default Swap valuation**  
Keywords: LIBOR market model, credit risk, Credit Default Swaption, Constant Maturity Credit Default Swap-method. (43 pages, 2006)
97. A. Dreyer  
**Interval Methods for Analog Circuits**  
Keywords: interval arithmetic, analog circuits, tolerance analysis, parametric linear systems, frequency response, symbolic analysis, CAD, computer algebra (36 pages, 2006)
98. N. Weigel, S. Weihe, G. Bitsch, K. Dreßler  
**Usage of Simulation for Design and Optimization of Testing**  
Keywords: Vehicle test rigs, MBS, control, hydraulics, testing philosophy (14 pages, 2006)
99. H. Lang, G. Bitsch, K. Dreßler, M. Speckert  
**Comparison of the solutions of the elastic and elastoplastic boundary value problems**  
Keywords: Elastic BVP, elastoplastic BVP, variational inequalities, rate-independency, hysteresis, linear kinematic hardening, stop- and play-operator (21 pages, 2006)
100. M. Speckert, K. Dreßler, H. Mauch  
**MBS Simulation of a hexapod based suspension test rig**  
Keywords: Test rig, MBS simulation, suspension, hydraulics, controlling, design optimization (12 pages, 2006)
101. S. Azizi Sultan, K.-H. Küfer  
**A dynamic algorithm for beam orientations in multicriteria IMRT planning**  
Keywords: radiotherapy planning, beam orientation optimization, dynamic approach, evolutionary algorithm, global optimization (14 pages, 2006)
102. T. Götz, A. Klar, N. Marheineke, R. Wegener  
**A Stochastic Model for the Fiber Lay-down Process in the Nonwoven Production**  
Keywords: fiber dynamics, stochastic Hamiltonian system, stochastic averaging (17 pages, 2006)
103. Ph. Süß, K.-H. Küfer  
**Balancing control and simplicity: a variable aggregation method in intensity modulated radiation therapy planning**  
Keywords: IMRT planning, variable aggregation, clustering methods (22 pages, 2006)
104. A. Beaudry, G. Laporte, T. Melo, S. Nickel  
**Dynamic transportation of patients in hospitals**  
Keywords: in-house hospital transportation, dial-a-ride, dynamic mode, tabu search (37 pages, 2006)
105. Th. Hanne  
**Applying multiobjective evolutionary algorithms in industrial projects**  
Keywords: multiobjective evolutionary algorithms, discrete optimization, continuous optimization, electronic circuit design, semi-infinite programming, scheduling (18 pages, 2006)
106. J. Franke, S. Halim  
**Wild bootstrap tests for comparing signals and images**  
Keywords: wild bootstrap test, texture classification, textile quality control, defect detection, kernel estimate, nonparametric regression (13 pages, 2007)
107. Z. Drezner, S. Nickel  
**Solving the ordered one-median problem in the plane**  
Keywords: planar location, global optimization, ordered median, big triangle small triangle method, bounds, numerical experiments (21 pages, 2007)
108. Th. Götz, A. Klar, A. Unterreiter, R. Wegener  
**Numerical evidence for the non-existing of solutions of the equations describing rotational fiber spinning**  
Keywords: rotational fiber spinning, viscous fibers, boundary value problem, existence of solutions (11 pages, 2007)
109. Ph. Süß, K.-H. Küfer  
**Smooth intensity maps and the Bortfeld-Boyer sequencer**  
Keywords: probabilistic analysis, intensity modulated radiotherapy treatment (IMRT), IMRT plan application, step-and-shoot sequencing (8 pages, 2007)
110. E. Ivanov, O. Gluchshenko, H. Andrä, A. Kudryavtsev  
**Parallel software tool for decomposing and meshing of 3d structures**  
Keywords: a-priori domain decomposition, unstructured grid, Delaunay mesh generation (14 pages, 2007)
111. O. Iliev, R. Lazarov, J. Willems  
**Numerical study of two-grid preconditioners for 1d elliptic problems with highly oscillating discontinuous coefficients**  
Keywords: two-grid algorithm, oscillating coefficients, preconditioner (20 pages, 2007)
112. L. Bonilla, T. Götz, A. Klar, N. Marheineke, R. Wegener  
**Hydrodynamic limit of the Fokker-Planck equation describing fiber lay-down processes**  
Keywords: stochastic differential equations, Fokker-Planck equation, asymptotic expansion, Ornstein-Uhlenbeck process (17 pages, 2007)
113. S. Rief  
**Modeling and simulation of the pressing section of a paper machine**  
Keywords: paper machine, computational fluid dynamics, porous media (41 pages, 2007)
114. R. Ciegis, O. Iliev, Z. Lakdawala  
**On parallel numerical algorithms for simulating industrial filtration problems**  
Keywords: Navier-Stokes-Brinkmann equations, finite volume discretization method, SIMPLE, parallel computing, data decomposition method (24 pages, 2007)
115. N. Marheineke, R. Wegener  
**Dynamics of curved viscous fibers with surface tension**  
Keywords: Slender body theory, curved viscous fibers with surface tension, free boundary value problem (25 pages, 2007)

116. S. Feth, J. Franke, M. Speckert

**Resampling-Methoden zur mse-Korrektur  
und Anwendungen in der Betriebsfestigkeit**

Keywords: Weibull, Bootstrap, Maximum-Likelihood,  
Betriebsfestigkeit

(16 pages, 2007)

117. H. Knaf

**Kernel Fisher discriminant functions – a concise and rigorous introduction**

Keywords: wild bootstrap test, texture classification,  
textile quality control, defect detection, kernel estimate,  
nonparametric regression

(30 pages, 2007)

118. O. Iliev, I. Rybak

**On numerical upscaling for flows in heterogeneous porous media**

Keywords: numerical upscaling, heterogeneous porous  
media, single phase flow, Darcy's law, multiscale problem,  
effective permeability, multipoint flux approximation,  
anisotropy

(17 pages, 2007)

119. O. Iliev, I. Rybak

**On approximation property of multipoint  
flux approximation method**

Keywords: Multipoint flux approximation, finite volume  
method, elliptic equation, discontinuous tensor coefficients,  
anisotropy

(15 pages, 2007)

Status quo: July 2007